



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Role of shoe cushioning, body mass and running biomechanics on injury risk: a study protocol for a randomised controlled trial

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-017379
Article Type:	Protocol
Date Submitted by the Author:	19-Apr-2017
Complete List of Authors:	Malisoux, Laurent; Luxembourg Institute of Health, Department of Population Health Delattre, Nicolas; Decathlon SportsLab, Movement Sciences Department Urhausen, Axel; Luxembourg Institute of Health, Department of Population Health; Centre Hospitalier de Luxembourg, Sports Clinic Theisen, Daniel; Luxembourg Institute of Health, Department of Population Health
Primary Subject Heading:	Sports and exercise medicine
Secondary Subject Heading:	Public health
Keywords:	Sports injury prevention, Footwear, Impact forces, EPIDEMIOLOGY

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TITLE PAGE

Title:

Role of shoe cushioning, body mass and running biomechanics on injury risk: a study protocol for a randomised controlled trial

Authors:

Laurent MALISOUX^a, Nicolas DELATTRE^b, Axel URHAUSEN^{a,c,d}, Daniel THEISEN^a

Affiliations:

^a Luxembourg Institute of Health, Sports Medicine Research Laboratory, Luxembourg, Grand-Duchy of Luxembourg

^b Decathlon SportsLab, Movement Sciences Department, Villeneuve d’Ascq, France

^c Centre Hospitalier de Luxembourg, Sports Clinic, Luxembourg, Grand-Duchy of Luxembourg

^d International University of Health, Exercise and Sports (LUNEX), 50 avenue du Parc des Sports L-4671 Differdange, Grand-Duchy of Luxembourg

Corresponding Author:

Laurent Malisoux
76 rue d’Eich
L-1460 Luxembourg
E-Mail: laurent.malisoux@lih.lu
Phone: +352 26970 231

Keywords:

Sports injury prevention, footwear, epidemiology, impact forces

Word count (excluding title page, abstract, references, figures and tables): 3567

Abstract Word count: 260

Number of figures: 1

Number of tables: 1

Online supplementary material: 3

Title: Role of shoe cushioning, body mass and running biomechanics on injury risk: a study protocol for a randomised controlled trial

ABSTRACT

Introduction: Repetitive loading of the musculoskeletal system is suggested to be involved in the underlying mechanism of the majority of running-related injuries (RRI). Accordingly, heavier runners are assumed to be at a higher risk of RRI. The cushioning system of modern running shoes is expected to protect runners against high impact forces, and therefore, RRI. However, the role of shoe cushioning in injury prevention remains unclear. The main aim of this study is to investigate the influence of shoe cushioning and body mass on RRI risk, while exploring simultaneously the association between running technique and RRI risk.

Methods and analysis: This double-blinded randomised controlled trial will involve about 800 healthy leisure-time runners. They will randomly receive one of two running shoe models that will differ in their cushioning properties (i.e. stiffness) by ~35%. The participants will perform a running test on an instrumented treadmill at their preferred running speed at baseline. They will then be followed-up prospectively over 6-month period, during which they will self-report all their sports activities as well as any injury in an internet-based database TIPPS (Training and Injury Prevention Platform for Sports). Cox regression analyses will be used to compare injury risk between the study groups, and to investigate the association between training, biomechanical and anatomical risk factors, and injury risk.

Ethics and dissemination: The study was approved by the National Ethics Committee for Research (Ref: 201701/02 v1.1). Outcomes will be disseminated through publications in peer-reviewed journals, presentations at international conferences, as well as articles in popular magazines and on specialised websites.

Trial registration number: NCT03115437

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66

55

56

57

58

59

60

61

62

63

64

65

66

STRENGTHS AND LIMITATIONS OF THIS STUDY

- Double-blinded randomised controlled trial (assessor and participant blinding) and intention-to-treat analysis.
- This study compares 2 shoe versions with widely differing cushioning properties while remaining within the cushioning range of models available on the market.
- A biomechanical analysis will be performed for each participant prior to the 6-month follow-up, which allows to investigate the association between running biomechanics and injury risk in a large cohort of runners.
- The running test will be carried out on a treadmill using a standardised protocol, which might not be reflective of the participants' habitual training conditions.

INTRODUCTION

Running is an increasingly popular form of physical activity. From a public health perspective, the promotion of leisure-time running might be a powerful strategy to combat the pandemic of physical inactivity worldwide,[1] and its consequence on non-communicable diseases.[2] Although regular running activity has a massive beneficial impact on health,[3] it also generates a relatively high number of injuries, especially at the lower limb.[4] The risk of sustaining a running-related injury (RRI) cancels out part of the benefits of running practice, since the long term consequences of injury include, among others, early-onset osteoarthritis,[5] a reduction of physical activity,[6] as well as an increase in health care costs.[7, 8] RRI incidence has remained high during the last 40 years, with an overall incidence rate ranging between 18.2% and 92.4%.[9] The role of footwear on RRI risk has been strongly emphasized ever since jogging became popular in the 1970s, but there is currently no evidence that developments in running shoe technology and new concepts regularly emerging on the market have helped to tackle the RRI burden.[10, 11]

Most RRI are overuse injuries, as they develop progressively over the kilometres run. The aetiology of these injuries is multifactorial,[12] which implies that to understand the mechanisms leading to injury, a holistic approach is warranted, including the study of a large set of potential risk factors. These factors could be classified as being related to training characteristics, running mechanics and anatomy of the runners. Some authors suggested that anatomical and biomechanical factors influence the tolerance to physical strain and thus the relationship between training load and injury occurrence.[13, 14]

Biological tissues such as bones, muscles and tendons can endure a certain amount of stress, provided that the product of stress level (e.g. intensity, external load) and the number of repetitions within a certain time period (e.g. strides, training sessions) remains below a threshold that is specific to each structure.[13] In running, the ground reaction force is the main external stress that acts on the body. Vertical ground reaction force (VGRF) is a biomechanical factor that has been extensively studied in

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

94 running.[15, 16] A recent meta-analysis found that the loading rate of the vertical ground reaction
95 force was higher in patients with a history of stress fracture.[15] High impact-related variables were
96 shown to increase the risk of bony and soft tissue injuries.[16] Moreover, running retraining
97 interventions have proven their efficiency in modifying some VGRF parameters and decreasing pain,
98 which suggest that running retraining represents an interesting paradigm to treat RRI.[17-19]

99

100 Since running biomechanics are associated with injury risk, any effect of shoe features on the running
101 pattern and VGRF parameters deserve attention. Given that repetitive loading of the musculoskeletal
102 system is an injury risk factor, cushioning has been one of the most extensively investigated shoe
103 feature. The shock absorption properties of footwear mainly result from the materials used in the sole
104 (i.e. their type, density, structure and combination), as well as from the geometry of the shoe (i.e. the
105 midsole thickness and the design of inserts). One of the most popular approaches has been to change
106 the hardness of the shoe midsole.[20-22] Overall, the studies investigating the effect of shoe
107 cushioning on VGRF did not provide consistent results. In theory, peak impact forces should be
108 reduced by softer or more compliant shoes,[23] which was indeed confirmed in some in vivo
109 studies.[24, 25] Conversely, some investigations did not find any effect of cushioning,[26] or reported
110 increased peak impact forces in softer shoes.[20, 27] Recently, a large cross sectional study revealed
111 that softer midsole hardness was associated with higher vertical force impact peak.[20] Unfortunately,
112 very few studies investigated the association between shoe cushioning and injury risk.[28, 29]
113 Therefore, the role of shoe cushioning systems in RRI prevention remains unknown.

114

115 Body mass index (BMI) has been associated with injury risk in novice,[30, 31] as well as in
116 recreational runners,[28] though other results suggest a protective effect of BMI.[9] It is common
117 belief that individuals with higher BMI have a higher injury risk, because of the increased physical
118 stress that results from extra body weight. Surprisingly, body mass as such has hardly ever been
119 considered as a potential risk factor for running injury.[9]

120

Surprisingly, the literature on the association between single shoe features and RRI risk is still poor.[11, 32, 33] Until now, no relationship has been found between the cushioning properties of modern running shoes and RRI risk,[28] but body mass should be taken into account here. Therefore, the main purpose of this study is to investigate the association between shoe cushioning and body mass on the one hand, and RRI risk on the other hand in recreational runners. The secondary aims are to identify which of the running technique-related characteristics (timing variables and VGRF parameters) are associated with injury risk, as well as with the cushioning properties of the shoes. The following hypotheses (H) will be tested:

H1. Running shoes with greater stiffness are associated with a higher injury risk in leisure-time runners.

H2. High body mass is associated with a higher injury risk in leisure-time runners.

H3. Runners with a high body mass experience a lower injury risk in shoes with greater stiffness.

H4. A higher step length, a lower step frequency, and higher peak vertical impact forces are associated with a higher injury risk.

H5. Running shoes with greater stiffness will be associated with higher vertical impact peak forces and a shorter contact time.

H6. High body mass will be associated with higher peak vertical impact forces, increased contact time, increased duty factor, and decreased step frequency.

Furthermore, exploratory risk factor analyses will be performed on the biomechanical variables obtained from the running analysis, anthropometric measurements, running experience, and habitual running speed. The focus of the analyses is the effect modification of body mass and other above mentioned risk factors on the association between shoe cushioning and injury risk.

143

144 **METHODS AND ANALYSIS**

145 **Trial design**

The design of this study is a randomised controlled trial with a 6-month follow-up and a biomechanical analysis of running pattern at baseline. The study is based on the comparison between

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2 running shoe prototypes, which only differ with respect to the cushioning (i.e. stiffness). The cushioning properties of both shoe versions are within the range of those from available models on the market. Running footwear is provided by a renowned sport equipment manufacturer. The main outcome is RRI (cf. definition below). The participants as well as the assessors are blinded to group allocation. The design of the trial is illustrated in Figure 1. The protocol conforms to the Recommendations for Interventional Trials (SPIRIT) and has been registered on <https://clinicaltrials.gov/> (NCT03115437, 11/04/2017).

Insert Figure 1 about here

Study population

The target population is leisure-time runners, regardless of running experience, fitness level, or body mass. Participants will be recruited through advertisements in local newspapers, social media, running magazines and press releases within the country during the months of September 2017 to January 2018. Healthy volunteers will be considered eligible if they are aged between 18 and 65 years and capable of performing 15 minutes of consecutive running. Volunteers will be excluded in case of any contraindication to perform running activity, prior (<12 months) surgery at the lower limbs or lower back region, use of orthopaedic insoles for running activities, or current RRI. Additionally, the participants will have to agree on the following requirements: 1) to practice running at least once a week, 2) to use the provided study shoes for all their running sessions, and 3) to report, at least once per week, all sports activities, as well as any injury or pain experienced during the follow-up period on an internet-based database called TIPPS (Training and Injury Prevention Platform for Sports, www.tipps.lu). Volunteers first have to create a personal account on TIPPS, pre-register to the study via their personal account, and answer an online inclusion/exclusion questionnaire as well as a baseline questionnaire. Answers to both questionnaires will be assessed by the investigators during the initial visit.

175 **Randomisation**

176 Participants must understand and agree on the randomized design of the study. Those who meet the
177 eligibility criteria and sign the informed consent form will be randomly allocated to one of the two
178 study arms. They will be stratified according to their sex, which is known to influence body mass as
179 well as many other anthropometric characteristics. Therefore, two pre-established randomisation lists
180 (block size = 40) will be prepared by a statistician not involved in any other part of the study before
181 the beginning of the recruitment. To ensure allocation concealment, the study groups and shoes will
182 be coded and the randomisation lists will be uploaded in the TIPPS system by an IT specialist who
183 will not be involved in any other part of the study. Then, the TIPPS system will provide the
184 investigator in charge of the recruitment with a study group number for each participant, according to
185 the randomisation lists. The investigator will upload the shoe number according to shoe size chosen
186 and study arm so that a cross validation will be performed by the electronic system. The investigators
187 in charge of the recruitment, the follow-up and data quality check, as well as the participants, will be
188 blinded regarding the shoe version distributed. The shoe code will be broken after completion of data
189 analysis.

191 **Intervention**

192 The study shoes are prototypes and will be anonymized for the purpose of this trial. The sole of the
193 shoes will be customized so that the two running shoe prototypes will be exactly the same (same
194 midsole, same outsole, same upper), except for their cushioning properties which will differ by about
195 35%, while remaining within the range of the models available on the market (stiffness: ~53-97
196 N/mm). The differences in cushioning properties between shoe versions will be created by modifying
197 the midsole material, i.e. chemistry, density, and therefore the hardness of the Ethylene Vinyl Acetate
198 (EVA) foam. In order to provide accurate data on the technical specifications (i.e. shoe stiffness) of
199 each prototype, a set of 40 shoes (10 pairs per condition) will be tested for cushioning properties by
200 the manufacturer according to a standardized protocol (Impact test: ASTM1614, Procedure A).[34]

201

Data collection

Baseline questionnaire

During the online registration process, the participants have to fill in a baseline questionnaire to report information regarding running experience, training habits, recent running competitions performed and injury history. A standardised questionnaire concerning the risk of sports participation must also be completed by the volunteers. Every participant responding positively to any of the symptom-based questions or to more than one risk factor will be invited for a clearance check by a sports medical doctor prior to the test.

Biomechanical testing

The biomechanical running analysis will be performed on an instrumented treadmill (M-Gait, Motekforce Link Amsterdam, The Netherlands) in the study shoes, according to the random allocation. The test (10 minutes) consists of a 5-minute warm-up followed by a 5-minute run at the self-declared preferred (habitual) running speed. Two records of 45 seconds will be obtained over the last 2 minutes of the test. No data will be recorded during the first 8 minutes, which was shown to be enough time to provoke short-term adaptations of running style with respect to the shoe type.[21, 35] Additionally, the participants who reported a preferred running speed equal to 10 km/h (+/- 1 km/h) will be invited to perform a second test at the end of the follow-up period. This second test will consist in 10 minutes of running in each shoe model. Records will be obtained during the last 2 minutes of each run. This will allow a within subject analysis of the shoe effect on running biomechanics at a standardised speed.

Table 1: Biomechanical variables of interest.

Variable	Abbreviation	Unit	Normalization
Step frequency	SF	[Steps.min ⁻¹]	/
Contact time	CT	[ms]	/
Flight time	FT	[ms]	/
Duty factor	DF	[%]	/

Step length	SL	[m]	[%LL]
Vertical Impact Peak Force	VIPF	[N]	[N.kg ⁻¹]
Peak Vertical Force	PVF	[N]	[N.kg ⁻¹]
Vertical Instantaneous Loading Rate	VILR	[N.s ⁻¹]	[N.kg ⁻¹ .s ⁻¹]
Vertical Average Loading Rate	VALR	[N.s ⁻¹]	[N.kg ⁻¹ .s ⁻¹]
Peak Power	PP	[W]	[W.kg ⁻¹]
Time to Peak Force	TPF	[ms]	/
Leg stiffness	Kleg	(kN/m)	/
Vertical stiffness	Kvert	(kN/m)	/

N: Newton, min: minute, ms: millisecond, m: meter, LL: leg length, kg: kilogram, W: Watt.

Anthropometric measures

The body mass of each participant will be measured before the treadmill running test in a stationary position. Also, the participants will have to report their body mass on a monthly basis onto their TIPPS account. Pop-up windows will inform the participants when an update is needed. In clinical settings, leg length is usually assessed as the measure between the anterior superior iliac spine and the medial malleolus, and is referred to as the “direct” clinical method.[36] The measurements will be performed on both legs and the average value will be used for the normalisation of step length. Additionally, the distance between the great trochanter and the ground will be measured to assess leg stiffness.[37] Body composition will be evaluated by bioelectrical impedance analysis (Tanita SC-240 MA). The proportion of fat mass will be included in the analyses as a potential confounder for the association between body mass and injury risk.

Data on exposure

Data on running practice will be collected using the TIPPS system.[28, 38] Required information in the sport activity report includes the type of activity, context, duration, subjectively perceived intensity, distance, shoe pair used, running surface (hard or soft), and whether the participant had

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

242 experienced any pain during the session forcing him/her to reduce practice volume or intensity, or to
243 interrupt the practice. Session intensity is determined using the Borg’s rating of perceived exertion
244 scale, a subjective 10-point scale.[39]

245
246 *Data on outcome*

247 The primary outcome is first-time RRI. A consensus definition of RRI in recreational runners has
248 been recently published.[40] The definition of RRI is a “running-related (training or competition)
249 musculoskeletal pain in the lower limbs that causes a restriction on or stoppage of running (distance,
250 speed, duration, or training) for at least 7 days or 3 consecutive scheduled training sessions, or that
251 requires the runner to consult a physician or other health professional.”
252 In previous studies, an RRI was defined as “any physical pain located at the lower limbs or lower
253 back region, sustained during or as a result of running practice and impeding planned running activity
254 for at least 1 day” (time-loss definition).[14, 28, 32, 33, 38] All injuries reported by the participants
255 during the follow-up will be assessed according to each of the two definitions presented here above.
256 The consensus definition will be considered as the reference, while a sensitive analysis will reveal if
257 the results would be impacted when using the former definition of RRI.

258 Similarly to uploading a training session or competition, the TIPPS provides a complete yet easy to
259 fill in questionnaire when reporting an injury. Information regarding the following is required: injury
260 date, context, sports discipline, injury mechanism (acute or progressive), anatomical location, type of
261 injury, description and estimated return date. RRI will be classified according to the Orchard Sports
262 Injury Classification System version 10 (OSICS-10).[41] Injury severity will be measured in days of
263 modified or interrupted training.

264
265 **Follow-up**

266 Given that the participants are required to practice running at least once a week, individual e-mail
267 reminders will be sent to the participants who do not provide the system with any data for the
268 previous week. Personal phone calls will be made if the participants do not react to the e-mail

reminders and if the reported information in either the training log or on the injury form is found to be inconsistent.

Injury data will be systematically checked by one of the investigators for completeness and coherence. Participants who do not complete their entire running calendar with weekly information will be contacted by one of the investigators to ensure that a RRI is not the reason for non-compliance or dropping out. The intervention period will last six months, allowing enough time for the participants to cover a large distance with the study shoes.

Sample size

A sample size calculation for Cox regression was used to determine the number of participants needed for the primary hypothesis of the study. With an alpha of 0.05 and a power of 80%, an average injury rate of 30%, [14, 32, 33] an expected HR=1.50 between groups, 50% of participants randomised to each shoe group and an expected drop-out rate of 20%, the total number of participants required is 802.

A within subject analysis will be performed on a subgroup of participants to investigate the effect of shoe condition on VGRF. A total sample of 39 participants will be required to detect a difference of 0.16 body weight (standard deviation: 0.25 body weight) [20] between shoe conditions with 80% power and a significant level of 5%.

Statistical analysis

Descriptive data for the personal, anthropometric, biomechanical and training-related characteristics will be presented as count and percentage for dichotomous variables, and as mean and standard deviation, or as median and range, respectively, for normally and non-normally distributed continuous variables. Average sport-related characteristics will be computed for each participant over their specific period of observation. Shock absorption properties of the two types of shoes will be compared using a Student's t test. A two-way analysis of variance (ANOVA) will be used to

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

determine whether any difference in running biomechanics results from the shoe cushioning properties or body mass.

Cox proportional hazards regressions will be used to compute the hazard rates (HR) in the exposure groups, using first-time injury as the primary outcome. Date of inclusion (baseline evaluation date) and date of injury or of censoring will be basic data used to calculate the time at risk, which is expressed in hours spent running and defined as the time-scale.[31] A participant will be right-censored if injury unrelated to running or severe disease caused a modification of the running plan, or at the end of follow-up. The assumption of proportional hazards will be evaluated by log-minus-log plots.

Unadjusted Cox regressions will be performed to present the crude estimates of HRs for shoe model, body mass and other potential risk factors such as running biomechanics variables and training-related characteristics. Body mass is an exposure that can change over time (time-dependent covariate). This means that each participant could move between exposure states continuously (every month in our study). A delayed entry will be used in the unadjusted Cox regression model for body mass.[42]

Subsequently, the variables with a P value <0.200 will be included in the adjusted Cox regression analysis to determine whether shoe cushioning and/or body mass are associated with injury risk, controlling for potential confounders. The recommendation for using at least 10 injuries per predictor variable included in the Cox regression analysis will be strictly followed.[43]

Finally, to investigate if the effect of shoe cushioning on RRI risk is modified by body mass, a stratified analysis will be performed using the median value of body mass as cut-off. HRs and their 95% confidence intervals (CI) will be determined within each stratum.[44] All analyses will be performed using STATA/SE version 14.

DISCUSSION

It is common belief that shoe cushioning technology protects the runner against harmful consequences of repetitive high-load impacts. Therefore, heavier runners are generally advised to use footwear with

adapted shock absorption properties. Surprisingly, few studies have investigated the impact of shoe cushioning on injury risk.[28, 29] These studies did not provide any evidence on the beneficial effect of increased shock absorption properties. However, none of them included anthropometric measures in their analyses. Also, one study compared different types of insoles added in the shoes,[29] while the other compared two versions of a standard running shoe with a limited difference in midsole hardness (~15%).[28] Other study limitations such as the sample size ($n < 250$)[28] or the study population (Royal Air Force recruits)[29] suggest that these results should be interpreted with caution. The evidence on the association between running shoe cushioning and RRI is still poor and inconclusive. One of the main reasons is the practical constraint of investigations trying to combine biomechanical analyses with a long-term prospective follow-up in a large number of runners.[11] This study is the first randomised controlled trial investigating the influence of shoe cushioning on RRI risk including an evaluation of running technique in all participants. The results will provide information on the real benefits provided by additional cushioning, as well as on the mechanisms that might explain any potential preventive effect.

ETHICS AND DISSEMINATION

This study will be conducted in accordance with the Declaration of Helsinki and the Medical Research Involving Human Subjects Act. Also, the study protocol was approved by the National Ethics Committee for Research (Ref: 201701/02 v1.1). Written informed consent will be obtained from all participants. All collected data will be stored electronically using a coding system. This will ensure that the data is used in the strictest confidence and will not reveal the identity of the participants. Collected raw data will not be passed on to unauthorised third parties. Results presented or published in articles and reports will be depicted in general terms, to maintain participant anonymity. Electronic data will be stored on a secure server in data files only accessible to the project leader and co-investigators of the project. A notification of this study was sent to the National Data Protection Agency (CNPd). Study results will be submitted for publication in peer-reviewed journals

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

and for presentation at international conferences. Furthermore, we aim to disseminate our results through popular specialised magazines and websites.

Contributors - LM, ND, AU and DT contributed to the study conception and study design. LM is the main investigator, wrote the article with input from other investigators, and will be responsible for the acquisition and analysis of the data. ND will be responsible for the shoe design, production and testing. LM and DT will be responsible for data interpretation and manuscript drafting. ND, AU and DT commented on the various versions of the study protocol. All authors approved the final manuscript.

Funding – This study is co-funded by Decathlon, Villeneuve d’Ascq, France and by the Ministry of Higher Education and Research of Luxembourg.

Competing interests – A research partnership agreement was signed between Decathlon and the LIH. ND is employed at Decathlon. Decathlon will not be involved in the collection, management, analysis and interpretation of data. LM, DT and AU may not gain or lose financially from the results of the study in any way.

Ethics approval - All procedures were approved by the National Ethics Committee for Research (Ref: 201701/02 v1.1).

370 **REFERENCE LIST**

- 371 1 Kohl HW, 3rd, Craig CL, Lambert EV, et al. The pandemic of physical inactivity: global
372 action for public health. *Lancet*. 2012;**380**:294-305.
- 373 2 World Health Organization (WHO). Global status report on noncommunicable diseases
374 2014. . *Geneva: World Health Organization (WHO)*. 2014.
- 375 3 Hespanhol Junior LC, Pillay JD, van Mechelen W, et al. Meta-Analyses of the Effects of
376 Habitual Running on Indices of Health in Physically Inactive Adults. *Sports Med*.
377 2015;**45**:1455-68.
- 378 4 Lopes AD, Hespanhol Junior LC, Yeung SS, et al. What are the main running-related
379 musculoskeletal injuries? A Systematic Review. *Sports Med*. 2012;**42**:891-905.
- 380 5 Leech RD, Edwards KL, Batt ME. Does running protect against knee osteoarthritis? Or
381 promote it? Assessing the current evidence. *Br J Sports Med*. 2015;**49**:1355-6.
- 382 6 Koplan JP, Rothenberg RB, Jones EL. The natural history of exercise: a 10-yr follow-up of
383 a cohort of runners. *Med Sci Sports Exerc*. 1995;**27**:1180-4.
- 384 7 Hespanhol Junior LC, van Mechelen W, Postuma E, et al. Health and economic burden of
385 running-related injuries in runners training for an event: A prospective cohort study. *Scand J*
386 *Med Sci Sports*. 2016;**26**:1091-9.
- 387 8 Hespanhol Junior LC, van Mechelen W, Verhagen E. Health and Economic Burden of
388 Running-Related Injuries in Dutch Trailrunners: A Prospective Cohort Study. *Sports Med*.
389 2016.
- 390 9 van Gent RN, Siem D, van Middelkoop M, et al. Incidence and determinants of lower
391 extremity running injuries in long distance runners: a systematic review. *Br J Sports Med*.
392 2007;**41**:469-80; discussion 80.
- 393 10 Knapik JJ, Jones BH, Steelman RA. Physical training in boots and running shoes: a
394 historical comparison of injury incidence in basic combat training. *Mil Med*. 2015;**180**:321-8.

1
2
3 395 11 Theisen D, Malisoux L, Gette P, et al. Footwear and running-related injuries –Running on
4
5 396 faith? *Sports Orthop Traumatol*. 2016;**32**:169-76.
6
7 397 12 Meeuwisse WH, Tyreman H, Hagel B, et al. A dynamic model of etiology in sport injury:
8
9 398 the recursive nature of risk and causation. *Clin J Sport Med*. 2007;**17**:215-9.
10
11 399 13 Hreljac A. Impact and overuse injuries in runners. *Med Sci Sports Exerc*. 2004;**36**:845-9.
12
13 400 14 Malisoux L, Nielsen RO, Urhausen A, et al. A step towards understanding the
14
15 401 mechanisms of running-related injuries. *J Sci Med Sport*. 2015;**18**:523-8.
16
17 402 15 van der Worp H, Vrielink JW, Bredeweg SW. Do runners who suffer injuries have higher
18
19 403 vertical ground reaction forces than those who remain injury-free? A systematic review and
20
21 404 meta-analysis. *Br J Sports Med*. 2016.
22
23 405 16 Davis IS, Bowser BJ, Mullineaux DR. Greater vertical impact loading in female runners
24
25 406 with medically diagnosed injuries: a prospective investigation. *Br J Sports Med*. 2015.
26
27 407 17 Barton CJ, Bonanno DR, Carr J, et al. Running retraining to treat lower limb injuries: a
28
29 408 mixed-methods study of current evidence synthesised with expert opinion. *Br J Sports Med*.
30
31 409 2016;**50**:513-26.
32
33 410 18 Diebal AR, Gregory R, Alitz C, et al. Forefoot running improves pain and disability
34
35 411 associated with chronic exertional compartment syndrome. *Am J Sports Med*. 2012;**40**:1060-
36
37 412 7.
38
39 413 19 Azevedo AP, Mezencio B, Amadio AC, et al. 16 Weeks of Progressive Barefoot Running
40
41 414 Training Changes Impact Force and Muscle Activation in Habitual Shod Runners. *PLoS One*.
42
43 415 2016;**11**:e0167234.
44
45 416 20 Baltich J, Maurer C, Nigg BM. Increased vertical impact forces and altered running
46
47 417 mechanics with softer midsole shoes. *PLoS One*. 2015;**10**:e0125196.
48
49 418 21 Hardin EC, van den Bogert AJ, Hamill J. Kinematic adaptations during running: effects of
50
51 419 footwear, surface, and duration. *Med Sci Sports Exerc*. 2004;**36**:838-44.
52
53
54
55
56
57
58
59
60

- 420 22 Nigg BM, Baltich J, Maurer C, et al. Shoe midsole hardness, sex and age effects on lower
421 extremity kinematics during running. *J Biomech.* 2012;**45**:1692-7.
- 422 23 Shorten M, Mientjes M. The 'heel impact' force peak during running is neither 'heel' nor
423 'impact' and does not quantify shoe cushioning effects. *Footwear Science.* 2011;**3**:41-58.
- 424 24 Logan S, Hunter I, JT JTH, et al. Ground reaction force differences between running
425 shoes, racing flats, and distance spikes in runners. *J Sports Sci Med.* 2010;**9**:147-53.
- 426 25 O'Leary K, Vorpahl KA, Heiderscheid B. Effect of cushioned insoles on impact forces
427 during running. *J Am Podiatr Med Assoc.* 2008;**98**:36-41.
- 428 26 Kersting UG, Bruggemann GP. Midsole material-related force control during heel-toe
429 running. *Res Sports Med.* 2006;**14**:1-17.
- 430 27 Nigg BM, Bahlisen HA, Luethi SM, et al. The influence of running velocity and midsole
431 hardness on external impact forces in heel-toe running. *J Biomech.* 1987;**20**:951-9.
- 432 28 Theisen D, Malisoux L, Genin J, et al. Influence of midsole hardness of standard
433 cushioned shoes on running-related injury risk. *Br J Sports Med.* 2014;**48**:371-6.
- 434 29 Withnall R, Eastaugh J, Freemantle N. Do shock absorbing insoles in recruits undertaking
435 high levels of physical activity reduce lower limb injury? A randomized controlled trial. *J R*
436 *Soc Med.* 2006;**99**:32-7.
- 437 30 Buist I, Bredeweg SW, Bessem B, et al. Incidence and risk factors of running-related
438 injuries during preparation for a 4-mile recreational running event. *Br J Sports Med.*
439 2010;**44**:598-604.
- 440 31 Nielsen RO, Malisoux L, Moller M, et al. Shedding Light on the Etiology of Sports
441 Injuries: A Look Behind the Scenes of Time-to-Event Analyses. *J Orthop Sports Phys Ther.*
442 2016:1-34.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

443 32 Malisoux L, Chambon N, Delattre N, et al. Injury risk in runners using standard or motion
444 control shoes: a randomised controlled trial with participant and assessor blinding. *Br J*
445 *Sports Med.* 2016;**50**:481-7.

446 33 Malisoux L, Chambon N, Urhausen A, et al. Influence of the Heel-to-Toe Drop of
447 Standard Cushioned Running Shoes on Injury Risk in Leisure-Time Runners: A Randomized
448 Controlled Trial With 6-Month Follow-up. *Am J Sports Med.* 2016.

449 34 International A. ASTM F1614-99 Standard test method for shock attenuating properties of
450 materials systems for athletic footwear. *ASTM International.* 2006:West Conshohocken, PA,
451 USA.

452 35 Delattre N, Chambon N, Berton E, et al. Effect of time during a running session with
453 minimal footwear. *Comput Methods Biomech Biomed Engin.* 2013;**16 Suppl 1**:104-5.

454 36 Sabharwal S, Kumar A. Methods for assessing leg length discrepancy. *Clin Orthop Relat*
455 *Res.* 2008;**466**:2910-22.

456 37 Morin JB, Dalleau G, Kyrolainen H, et al. A simple method for measuring stiffness during
457 running. *J Appl Biomech.* 2005;**21**:167-80.

458 38 Malisoux L, Ramesh J, Mann R, et al. Can parallel use of different running shoes decrease
459 running-related injury risk? *Scand J Med Sci Sports.* 2015;**25**:110-5.

460 39 Foster C. Monitoring training in athletes with reference to overtraining syndrome. *Med Sci*
461 *Sports Exerc.* 1998;**30**:1164-8.

462 40 Yamato TP, Saragiotto BT, Lopes AD. A consensus definition of running-related injury in
463 recreational runners: a modified Delphi approach. *J Orthop Sports Phys Ther.* 2015;**45**:375-
464 80.

465 41 Rae K, Orchard J. The Orchard Sports Injury Classification System (OSICS) version 10.
466 *Clin J Sport Med.* 2007;**17**:201-4.

- 467 42 Nielsen RO, Parner ET, Nohr EA, et al. Excessive progression in weekly running distance
468 and risk of running-related injuries: an association which varies according to type of injury. *J*
469 *Orthop Sports Phys Ther.* 2014;**44**:739-47.
- 470 43 Peduzzi P, Concato J, Feinstein AR, et al. Importance of events per independent variable
471 in proportional hazards regression analysis. II. Accuracy and precision of regression
472 estimates. *J Clin Epidemiol.* 1995;**48**:1503-10.
- 473 44 Knol MJ, VanderWeele TJ. Recommendations for presenting analyses of effect
474 modification and interaction. *Int J Epidemiol.* 2012;**41**:514-20.

475

476

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE LEGEND


Figure 1: Trial design.

SUPPLEMENTARY FILES

Supplementary file 1: SPIRIT Checklist

Supplementary file 1: Study schedule

Figure 1:





SPIRIT 2013 Checklist: Recommended items to address in a clinical trial protocol and related documents*

Section/item	Item No	Description	Addressed on page number
Administrative information			
Title	1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	Page 1
Trial registration	2a	Trial identifier and registry name. If not yet registered, name of intended registry	Page 5: clinicaltrials.gov (NCT03115437)
	2b	All items from the World Health Organization Trial Registration Data Set	/
Protocol version	3	Date and version identifier	Page 7
Funding	4	Sources and types of financial, material, and other support	Page 15
Roles and responsibilities	5a	Names, affiliations, and roles of protocol contributors	Page 15
	5b	Name and contact information for the trial sponsor	Page 15
	5c	Role of study sponsor and funders, if any, in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication, including whether they will have ultimate authority over any of these activities	Page 15

5d	Composition, roles, and responsibilities of the coordinating centre, steering committee, endpoint adjudication committee, data management team, and other individuals or groups overseeing the trial, if applicable (see Item 21a for data monitoring committee)	n.a.
----	--	------

Introduction

Background and rationale	6a	Description of research question and justification for undertaking the trial, including summary of relevant studies (published and unpublished) examining benefits and harms for each intervention	Pages 4 and 5
	6b	Explanation for choice of comparators	Pages 5 and 4
Objectives	7	Specific objectives or hypotheses	Page 6
Trial design	8	Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group), allocation ratio, and framework (eg, superiority, equivalence, noninferiority, exploratory)	Page 6 and 7

Methods: Participants, interventions, and outcomes

Study setting	9	Description of study settings (eg, community clinic, academic hospital) and list of countries where data will be collected. Reference to where list of study sites can be obtained	Page 7
Eligibility criteria	10	Inclusion and exclusion criteria for participants. If applicable, eligibility criteria for study centres and individuals who will perform the interventions (eg, surgeons, psychotherapists)	Page 7
Interventions	11a	Interventions for each group with sufficient detail to allow replication, including how and when they will be administered	Page 8
	11b	Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose change in response to harms, participant request, or improving/worsening disease)	Page 11-12
	11c	Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence (eg, drug tablet return, laboratory tests)	Page 11-12
	11d	Relevant concomitant care and interventions that are permitted or prohibited during the trial	n.a.

1				
2				
3	Outcomes	12	Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg, median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen efficacy and harm outcomes is strongly recommended	Page 9 and 11
4				
5				
6				
7				
8	Participant timeline	13	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for participants. A schematic diagram is highly recommended (see Figure)	Figure 1 and Suppl. file 2
9				
10				
11	Sample size	14	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	Page 12
12				
13				
14	Recruitment	15	Strategies for achieving adequate participant enrolment to reach target sample size	Page 7
15				
16				

17 **Methods: Assignment of interventions (for controlled trials)**

18 Allocation:

19				
20				
21	Sequence generation	16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	Page 8
22				
23				
24				
25				
26	Allocation concealment mechanism	16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	Page 8
27				
28				
29				
30	Implementation	16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	Page 8
31				
32				
33				
34	Blinding (masking)	17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	Page 8
35				
36				
37		17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	Page 8
38				
39				

40 **Methods: Data collection, management, and analysis**

Data collection methods	18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known. Reference to where data collection forms can be found, if not in the protocol	Pages 9 to 11
	18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols	Page 11
Data management	19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol	Page 12
Statistical methods	20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol	Pages 12 and 13
	20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	Pages 12 and 13
	20c	Definition of analysis population relating to protocol non-adherence (eg, as randomised analysis), and any statistical methods to handle missing data (eg, multiple imputation)	Pages 12 and 13
Methods: Monitoring			
Data monitoring	21a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of whether it is independent from the sponsor and competing interests; and reference to where further details about its charter can be found, if not in the protocol. Alternatively, an explanation of why a DMC is not needed	/
	21b	Description of any interim analyses and stopping guidelines, including who will have access to these interim results and make the final decision to terminate the trial	/
Harms	22	Plans for collecting, assessing, reporting, and managing solicited and spontaneously reported adverse events and other unintended effects of trial interventions or trial conduct	Page 12
Auditing	23	Frequency and procedures for auditing trial conduct, if any, and whether the process will be independent from investigators and the sponsor	/

Ethics and dissemination


1				
2				
3	Research ethics	24	Plans for seeking research ethics committee/institutional review board (REC/IRB) approval	Page 14
4	approval			
5				
6	Protocol	25	Plans for communicating important protocol modifications (eg, changes to eligibility criteria, outcomes,	Page 14
7	amendments		analyses) to relevant parties (eg, investigators, REC/IRBs, trial participants, trial registries, journals,	
8			regulators)	
9				
10	Consent or assent	26a	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and	Page 7
11			how (see Item 32)	
12				
13		26b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary	n.a.
14			studies, if applicable	
15				
16	Confidentiality	27	How personal information about potential and enrolled participants will be collected, shared, and maintained	Page 7
17			in order to protect confidentiality before, during, and after the trial	
18				
19	Declaration of	28	Financial and other competing interests for principal investigators for the overall trial and each study site	Page 15
20	interests			
21				
22	Access to data	29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that	Pages 14 and 15
23			limit such access for investigators	
24				
25	Ancillary and post-	30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial	/
26	trial care		participation	
27				
28	Dissemination policy	31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals,	Page 14
29			the public, and other relevant groups (eg, via publication, reporting in results databases, or other data	
30			sharing arrangements), including any publication restrictions	
31				
32		31b	Authorship eligibility guidelines and any intended use of professional writers	/
33				
34		31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	n.a.
35				
36				
37	Appendices			
38				
39	Informed consent	32	Model consent form and other related documentation given to participants and authorised surrogates	Supplementary file
40	materials			3
41				
42				
43				
44				
45				
46				
47				
48				
49				

Biological specimens	33	Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in the current trial and for future use in ancillary studies, if applicable	n.a.
----------------------	----	--	------

*It is strongly recommended that this checklist be read in conjunction with the SPIRIT 2013 Explanation & Elaboration for important clarification on the items. Amendments to the protocol should be tracked and dated. The SPIRIT checklist is copyrighted by the SPIRIT Group under the Creative Commons "[Attribution-NonCommercial-NoDerivs 3.0 Unported](#)" license.

For peer review only

Supplementary file 2: Schedule of enrolment, interventions, and assessments for the study

	STUDY PERIOD							
	Enrolment	Allocation	Post-allocation: For every sport activity					Close-out
TIMEPOINT	-t ₁	0	t ₁	t ₂	t ₃	t ₄	etc.	6 Months
ENROLMENT								
Eligibility screen	X							
Informed consent	X							
Baseline Questionnaire	X	X						
Allocation		X						
Shoe distribution		X						
Running analysis		X						
INTERVENTIONS								
[Intervention A]								
[Intervention B]								
ASSESSMENTS								
Running experience	X	X						
Running regularity	X	X						
Typical running frequency	X	X						
Typical running distance	X	X						
Training running speed	X	X						
Type of running	X	X						
Competition participation	X	X						
Last event distance	X	X						
Favourite running distance	X	X						
Best time on 5 km / 10km	X	X						
Previous injury	X	X						

Height	X	X						
Body mass	X	X						
Leg length	X	X						
% fat tissue	X	X						
Step frequency	X	X						
Contact time	X	X						
Flight time	X	X						
Duty factor	X	X						
Step length	X	X						
Vertical Impact Peak Force	X	X						
Peak Vertical Force	X	X						
Vertical Instantaneous Loading Rate	X	X						
Vertical Average Loading Rate	X	X						
Peak Power	X	X						
Time to Peak Force	X	X						
Leg stiffness	X	X						
Vertical stiffness	X	X						
Sports discipline			X	X	X	X	etc.	
Duration			X	X	X	X	etc.	
Distance (if applicable)			X	X	X	X	etc.	
Perceived Exertion			X	X	X	X	etc.	
Shoe used (if running)			X	X	X	X	etc.	
Surface (if running)			X	X	X	X	etc.	
Pain*			X	X	X	X	etc.	
Injury**			X	X	X	X	etc.	

*The pain did not stop the participant from continuing normal training

**The participants had to adapt or interrupt their training accordingly

Free Informed Consent

Title:
Institution:
Project manager:
Research assistant:
Head of unit:

1. I declare to have read the above-described information and accept to voluntarily participate in the study “Effects of bodyweight and shoe cushioning on injury risk and running biomechanics: A randomised control trial” conducted by the SMRL.
2. I accept that my data shall be used and communicated to the commercial partner for strictly scientific purposes once it has been pseudonymised (coded).
3. I received a copy of the present signed informed consent document, as well as the general information intended for athlete participants. I received a clear description of the purpose and the nature of the study and I am aware of what is expected of me as a participant in this study. I have had enough time and the opportunity to ask questions about the study; all my questions have been met with a satisfactory answer.
4. I am free to retire from the study at any time without justification. By doing so I will not suffer any material or moral damage.
5. I agree that the results of this study can be subject to public talks or scientific publication.
6. I voluntarily consent to participate in this study and I fully understand what kind of data will be gathered during the study.
7. I preserve/abide the rights of access, deletion or modification of my personal data. Any personal information will be kept confidential and protected in agreement with the modified personal data protection act of August 2nd 2002. I can exercise that right via the project manager.

The responding signatory freely consents to participate in the above mentioned study

Name and First Name of the respondent:

Signature of the respondent:

Name and signature of the project manager:

Place and date:

BMJ Open

Shoe cushioning, body mass and running biomechanics as risk factors for running injury: a study protocol for a randomised controlled trial

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-017379.R1
Article Type:	Protocol
Date Submitted by the Author:	02-Jun-2017
Complete List of Authors:	Malisoux, Laurent; Luxembourg Institute of Health, Department of Population Health Delattre, Nicolas; Decathlon SportsLab, Movement Sciences Department Urhausen, Axel; Luxembourg Institute of Health, Department of Population Health; Centre Hospitalier de Luxembourg, Sports Clinic Theisen, Daniel; Luxembourg Institute of Health, Department of Population Health
Primary Subject Heading:	Sports and exercise medicine
Secondary Subject Heading:	Public health
Keywords:	Sports injury prevention, Footwear, Impact forces, EPIDEMIOLOGY

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TITLE PAGE

Title:

Shoe cushioning, body mass and running biomechanics as risk factors for running injury: a study protocol for a randomised controlled trial

Authors:

Laurent MALISOUX^a, Nicolas DELATTRE^b, Axel URHAUSEN^{a,c,d}, Daniel THEISEN^a

Affiliations:

^a Luxembourg Institute of Health, Sports Medicine Research Laboratory, Luxembourg, Grand-Duchy of Luxembourg

^b Decathlon SportsLab, Movement Sciences Department, Villeneuve d’Ascq, France

^c Centre Hospitalier de Luxembourg, Sports Clinic, Luxembourg, Grand-Duchy of Luxembourg

^d International University of Health, Exercise and Sports (LUNEX), 50 avenue du Parc des Sports L-4671 Differdange, Grand-Duchy of Luxembourg

Corresponding Author:

Laurent Malisoux
76 rue d’Eich
L-1460 Luxembourg
E-Mail: laurent.malisoux@lih.lu
Phone: +352 26970 231

Keywords:

Sports injury prevention, footwear, epidemiology, impact forces

Word count (excluding title page, abstract, references, figures and tables): 3472

Abstract Word count: 260

Number of figures: 1

Number of tables: 1

Online supplementary material: 4

Title: Shoe cushioning, body mass and running biomechanics as risk factors for running injury: a study protocol for a randomised controlled trial

ABSTRACT

Introduction: Repetitive loading of the musculoskeletal system is suggested to be involved in the underlying mechanism of the majority of running-related injuries (RRI). Accordingly, heavier runners are assumed to be at a higher risk of RRI. The cushioning system of modern running shoes is expected to protect runners against high impact forces, and therefore, RRI. However, the role of shoe cushioning in injury prevention remains unclear. The main aim of this study is to investigate the influence of shoe cushioning and body mass on RRI risk, while exploring simultaneously the association between running technique and RRI risk.

Methods and analysis: This double-blinded randomised controlled trial will involve about 800 healthy leisure-time runners. They will randomly receive one of two running shoe models that will differ in their cushioning properties (i.e. stiffness) by ~35%. The participants will perform a running test on an instrumented treadmill at their preferred running speed at baseline. They will then be followed-up prospectively over a 6-month period, during which they will self-report all their sports activities as well as any injury in an internet-based database TIPPS (Training and Injury Prevention Platform for Sports). Cox regression analyses will be used to compare injury risk between the study groups and to investigate the association between training, biomechanical and anatomical risk factors, and injury risk.

Ethics and dissemination: The study was approved by the National Ethics Committee for Research (Ref: 201701/02 v1.1). Outcomes will be disseminated through publications in peer-reviewed journals, presentations at international conferences, as well as articles in popular magazines and on specialised websites.

Trial registration number: NCT03115437

1
2
3 55
4
5 56
6
7 57
8
9 58
10
11 59
12
13 60
14
15 61
16
17 62
18
19 63
20
21 64
22
23 65
24
25 66
26
27 67
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STRENGTHS AND LIMITATIONS OF THIS STUDY

- Double-blinded randomised controlled trial (assessor and participant blinding) and intention-to-treat analysis.
- This study compares 2 shoe versions with widely differing cushioning properties while remaining within the cushioning range of models available on the market.
- A biomechanical analysis will be performed for each participant prior to the 6-month follow-up, which allows to investigate the association between running biomechanics and injury risk in a large cohort of runners.
- The running test will be carried out on a treadmill using a standardised protocol, which might not be reflective of the participants' habitual training conditions.

INTRODUCTION

Running is an increasingly popular form of physical activity. From a public health perspective, the promotion of leisure-time running might be a powerful strategy to combat the pandemic of physical inactivity worldwide,[1] and its consequence on non-communicable diseases.[2] Although regular running activity has a massive beneficial impact on health,[3] it also generates a relatively high number of injuries, especially at the lower limb.[4] The risk of sustaining a running-related injury (RRI) cancels out part of the benefits of running practice, since the long term consequences of injury might include, among others, increased risk of osteoarthritis,[5] a reduction of physical activity,[6] as well as an increase in health care costs.[7, 8] RRI incidence has remained high during the last 40 years, with an overall incidence rate ranging between 18.2% and 92.4%.[9] The role of footwear on RRI risk has been strongly emphasized ever since jogging became popular in the 1970s, but there is currently no evidence that developments in running shoe technology and new concepts regularly emerging on the market have helped to tackle the RRI burden.[10-12]

Most RRI are overuse injuries, as they develop progressively over the kilometres run. The aetiology of these injuries is multifactorial,[13] which implies that to understand the mechanisms leading to injury, a holistic approach is warranted, including the study of a large set of potential risk factors. These factors could be classified as being related to training characteristics, running mechanics and anatomy of the runners. Some authors suggested that anatomical and biomechanical factors influence the tolerance to physical strain and thus the relationship between training load and injury occurrence.[14, 15]

Biological tissues such as bones, muscles and tendons can endure a certain amount of stress, provided that the product of stress level (e.g. intensity, external load) and the number of repetitions within a certain time period (e.g. strides, training sessions) remains below a threshold that is specific to each structure.[14] In running, the ground reaction force is the main external stress that acts on the body. Vertical ground reaction force (VGRF) is a biomechanical factor that has been extensively studied in

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

95 running.[16, 17] A recent meta-analysis found that the loading rate of the vertical ground reaction
96 force was higher in patients with a history of stress fracture.[16] High impact-related variables were
97 shown to increase the risk of bony and soft tissue injuries.[17] Moreover, running retraining
98 interventions have proven their efficiency in modifying some VGRF parameters and decreasing pain,
99 which suggest that running retraining represents an interesting paradigm to treat RRI.[18-20] Other
100 biomechanical factors such as step length,[21] step frequency [22] or leg stiffness [23] have
101 previously been suggested as potential biomechanical risk factors for RRI, yet no causal relationship
102 has been established.
103
104 Since running biomechanics are associated with injury risk, any effect of shoe features on the running
105 pattern and VGRF parameters deserve attention. Given that repetitive loading of the musculoskeletal
106 system is an injury risk factor, cushioning has been one of the most extensively investigated shoe
107 features. The shock absorption properties of footwear mainly result from the materials used in the sole
108 (i.e. their type, density, structure and combination), as well as from the geometry of the shoe (i.e. the
109 midsole thickness and the design of inserts). One of the most popular approaches has been to change
110 the hardness of the shoe midsole.[24-26] Overall, the studies investigating the effect of shoe
111 cushioning on VGRF did not provide consistent results. In theory, peak impact forces should be
112 reduced by softer or more compliant shoes,[27] which was indeed confirmed in some in vivo
113 studies.[28, 29] Conversely, some investigations did not find any effect of cushioning,[30] or reported
114 increased peak impact forces in softer shoes.[24, 31] Recently, a large cross sectional study revealed
115 that softer midsole hardness was associated with higher vertical force impact peak.[24] Unfortunately,
116 very few studies have investigated the association between shoe cushioning and injury risk.[32, 33] In
117 a previous randomised controlled trial, midsole hardness was not associated with RRI risk. However,
118 the difference in shoe stiffness between the shoe conditions was limited (15%).[32] Therefore, the
119 role of shoe cushioning systems in RRI prevention remains unclear.

1
2
3 121 Body mass index (BMI) has been associated with injury risk in novice,[34, 35] as well as in
4
5 122 recreational runners,[32] though other results suggest a protective effect of BMI.[9] It is common
6
7 123 belief that individuals with higher BMI have a higher injury risk, because of the increased physical
8
9 124 stress that results from extra body weight. Surprisingly, body mass as such has hardly ever been
10
11 125 considered as a potential risk factor for running injury.[9]
12

13
14 126

15 127 Surprisingly, the literature on the association between single shoe features and RRI risk is still
16
17 128 poor.[11, 36, 37] Until now, no relationship has been found between the cushioning properties of
18
19 129 modern running shoes and RRI risk,[32] but body mass should be taken into account here. Therefore,
20
21 130 the main purpose of this study is to investigate the association between shoe cushioning and body
22
23 131 mass on the one hand, and RRI risk on the other hand in recreational runners. The secondary aims are
24
25 132 to identify which of the running technique-related characteristics (timing variables and VGRF
26
27 133 parameters) are associated with injury risk, as well as with the cushioning properties of the shoes.
28
29 134 Shoe cushioning will be characterised by the stiffness at the heel (N/mm) and quantified by
30
31 135 standardised impact test.[38] The following hypotheses (H) will be tested:

32
33 136 H1. Running shoes with greater stiffness are associated with a higher injury risk in leisure-time
34
35 137 runners.

36
37 138 H2. High body mass is associated with a higher injury risk in leisure-time runners.

38
39 139 H3. Runners with a high body mass experience a lower injury risk in shoes with greater stiffness.

40
41 140 H4. A higher step length, a lower step frequency, and higher vertical loading rate are associated with a
42
43 141 higher injury risk.

44
45 142 H5. Running shoes with greater stiffness will be associated with higher vertical loading rate and a
46
47 143 shorter contact time.

48
49 144 H6. High body mass will be associated with higher vertical loading rate, increased contact time,
50
51 145 increased duty factor, and decreased step frequency.

52
53 146 Furthermore, exploratory risk factor analyses will be performed on the biomechanical variables
54
55 147 obtained from the running analysis, anthropometric measurements, running experience, and habitual
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

running speed. The focus of the analyses is the effect modification of body mass and other above mentioned risk factors on the association between shoe cushioning and injury risk.

METHODS AND ANALYSIS

Trial design

The design of this study is a randomised controlled trial with a 6-month follow-up and a biomechanical analysis of running pattern at baseline. The study is based on the comparison between 2 running shoe prototypes, which only differ with respect to the cushioning (i.e. stiffness). The cushioning properties of both shoe versions are within the range of those from available models on the market. Running footwear is provided by a renowned sport equipment manufacturer. The main outcome is RRI (cf. definition below). The participants as well as the assessors are blinded to group allocation. The design of the trial is illustrated in Figure 1. The protocol conforms to the Recommendations for Interventional Trials (SPIRIT, supplementary files 1 and 2) and has been registered on <https://clinicaltrials.gov/> (NCT03115437, 11/04/2017).

Insert Figure 1 about here

Study population

The target population is leisure-time runners, regardless of running experience, fitness level, or body mass. Participants will be recruited through advertisements in local newspapers, social media, running magazines and press releases within the country during the months of September 2017 to January 2018. Healthy volunteers will be considered eligible if they are aged between 18 and 65 years and capable of performing 15 minutes of consecutive running. Volunteers will be excluded in case of any contraindication to perform running activity, prior (<12 months) surgery or major trauma to the lower limbs or lower back region, any running impeding injury over the previous months, or use of orthopaedic insoles for running activities. Additionally, the participants will have to agree on the following requirements: 1) to practice running at least once a week, 2) to use the provided study shoes

for all their running sessions, and 3) to report, at least once per week, all sports activities, as well as any injury or pain experienced during the follow-up period on an internet-based database called TIPPS (Training and Injury Prevention Platform for Sports, www.tipps.lu). Volunteers first have to create a personal account on TIPPS, pre-register to the study via their personal account, and answer an online inclusion/exclusion questionnaire as well as a baseline questionnaire. Answers to both questionnaires will be assessed by the investigators during the initial visit.

181

182 **Randomisation**

Participants must understand and agree on the randomized design of the study. Those who meet the eligibility criteria and sign the informed consent form will be randomly allocated to one of the two study arms. They will be stratified according to their sex, which is known to influence body mass as well as many other anthropometric characteristics. Therefore, two pre-established randomisation lists (block size = 40) will be prepared by a statistician not involved in any other part of the study before the beginning of the recruitment. To ensure allocation concealment, the study groups and shoes will be coded and the randomisation lists will be uploaded in the TIPPS system by an IT specialist who will not be involved in any other part of the study. Then, the TIPPS system will provide the investigator in charge of the recruitment with a study group number for each participant, according to the randomisation lists. The investigator will upload the shoe number according to shoe size chosen and study arm so that a cross validation will be performed by the electronic system. The investigators in charge of the recruitment, the follow-up and data quality check, as well as the participants, will be blinded regarding the shoe version distributed. The shoe code will be broken after completion of data analysis.

197

198 **Intervention**

The study shoes are prototypes and will be anonymized for the purpose of this trial. The sole of the shoes will be customized so that the two running shoe prototypes will be exactly the same (same midsole, same outsole, same upper), except for their cushioning properties which will differ by about

35%, while remaining within the range of the models available on the market (stiffness: ~53-97 N/mm). The differences in cushioning properties between shoe versions will be created by modifying the midsole material, i.e. chemistry, density, and therefore the hardness of the Ethylene Vinyl Acetate (EVA) foam. In order to provide accurate data on the technical specifications (i.e. shoe stiffness) of each prototype, a set of 40 shoes (10 pairs per condition) will be tested for stiffness properties by the manufacturer according to a standardized protocol (Impact test: ASTM1614, Procedure A).[38]

Data collection

Baseline questionnaire

During the online registration process, the participants have to fill in a baseline questionnaire to report information regarding running experience, training habits, recent running competitions performed and injury history. A standardised questionnaire concerning the risk of sports participation must also be completed by the volunteers (Supplementary file 3). Every participant responding positively to any of the symptom-based questions or presenting more than one cardiovascular risk factor will be invited for a clearance check by a sports medical doctor prior to the test.

Biomechanical testing

The biomechanical running analysis will be performed on an instrumented treadmill (M-Gait, Motekforce Link Amsterdam, The Netherlands) in the randomly allocated study shoes. The test (10 minutes) consists of a 5-minute warm-up followed by a 5-minute run at the self-declared preferred (habitual) running speed. Two records of 45 seconds will be obtained at a sampling rate of 1 kHz over the last 2 minutes of the test. No data will be recorded during the first 8 minutes, which was shown to be enough time to provoke short-term adaptations of running style with respect to the shoe type.[25, 39] The main biomechanical variables of interest are presented in table 1.

Table 1: Biomechanical variables of interest.

Variable	Abbreviation	Unit	Normalization
----------	--------------	------	---------------

Step frequency	SF	[Steps.min ⁻¹]	/
Contact time	CT	[ms]	/
Flight time	FT	[ms]	/
Duty factor	DF	[%]	/
Step length	SL	[m]	[%LL]
Vertical Impact Peak Force	VIPF	[N]	[N.kg ⁻¹]
Peak Vertical Force	PVF	[N]	[N.kg ⁻¹]
Vertical Instantaneous Loading Rate	VILR	[N.s ⁻¹]	[N.kg ⁻¹ .s ⁻¹]
Vertical Average Loading Rate	VALR	[N.s ⁻¹]	[N.kg ⁻¹ .s ⁻¹]
Peak Power	PP	[W]	[W.kg ⁻¹]
Time to Peak Force	TPF	[ms]	/
Leg stiffness	Kleg	(kN/m)	/
Vertical stiffness	Kvert	(kN/m)	/

N: Newton, min: minute, ms: millisecond, m: meter, LL: leg length, kg: kilogram, W: Watt.

Anthropometric measures

Body mass and height of each participant will be measured barefoot and in running clothes before the treadmill running test. Also, the participants will have to report their body mass on a monthly basis onto their TIPPS account. Pop-up windows will inform the participants when an update is needed. In clinical settings, leg length is usually assessed as the measure between the anterior superior iliac spine and the medial malleolus (supine position), and is referred to as the “direct” clinical method.[40] The measurements will be performed on both legs and the average value will be used for the normalisation of step length. Additionally, the distance between the greater trochanter and the ground will be measured (standing position) to assess leg stiffness.[41] Body composition will be evaluated by bioelectrical impedance analysis (Tanita SC-240 MA). The proportion of fat mass will be included in the analyses as a potential confounder for the association between body mass and injury risk.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

242 *Data on exposure*

243 Data on running practice will be collected using the TIPPS system.[32, 42] Required information in
244 the sport activity report includes the type of activity, context, duration, subjectively perceived
245 intensity, distance, shoe pair used, running surface (hard or soft), and whether the participant had
246 experienced any pain during the session forcing him/her to reduce practice volume or intensity, or to
247 interrupt the practice. Session intensity is determined using the Borg’s rating of perceived exertion
248 scale, a subjective 10-point scale.[43]

250 *Data on outcome*

251 The primary outcome is first-time RRI. A consensus definition of RRI in recreational runners has
252 been recently published.[44] The definition of RRI is a “running-related (training or competition)
253 musculoskeletal pain in the lower limbs that causes a restriction on or stoppage of running (distance,
254 speed, duration, or training) for at least 7 days or 3 consecutive scheduled training sessions, or that
255 requires the runner to consult a physician or other health professional.”
256 In previous studies, an RRI was defined as “any physical pain located at the lower limbs or lower
257 back region, sustained during or as a result of running practice and impeding planned running activity
258 for at least 1 day” (time-loss definition).[15, 32, 36, 37, 42] All painful episodes reported by the
259 participants during the follow-up will be assessed by a member of the research team according to each
260 of the two definitions presented above. The consensus definition will be considered as the reference,
261 while a sensitivity analysis will reveal if the results would be impacted when using the former
262 definition of RRI.
263 Similarly to uploading a training session or competition, the TIPPS provides a complete yet easy to
264 fill in questionnaire when reporting an injury. Information regarding the following is required: injury
265 date, context, sports discipline, injury mechanism (acute or progressive), anatomical location, type of
266 injury, description (free text field) and estimated return date. RRIs will be classified according to the
267 Orchard Sports Injury Classification System version 10 (OSICS-10).[45] Injury severity will be
268 measured in days of modified or interrupted training.

269

270 Follow-up

271 Given that the participants are required to practice running at least once a week, individual e-mail
272 reminders will be sent to the participants who do not provide the system with any data for the
273 preceding week. Personal phone calls will be made if the participants do not react to the e-mail
274 reminders and if the reported information in either the training log or on the injury form is found to be
275 inconsistent.

276 Participants reporting any injury will be systematically contacted by one of the investigators to verify
277 completeness and coherence of the reported data, and to check if the episode qualifies as an RRI (as
278 defined above). Participants who do not complete their entire running calendar with weekly
279 information will be contacted by one of the investigators to ensure that a RRI is not the reason for
280 non-compliance or dropping out. The intervention period will last six months, allowing enough time
281 for the participants to cover a large distance with the study shoes.

282

283 Sample size

284 A sample size calculation for Cox regression was used to determine the number of participants needed
285 for the primary hypothesis of the study. With an alpha of 0.05 and a power of 80%, an average injury
286 rate of 30%, [15, 36, 37] an expected hazard rate ratio (HR)=1.50 between groups, 50% of participants
287 randomised to each shoe group and an expected drop-out rate of 20%, the total number of participants
288 required is 802.

289

290 Statistical analysis

291 Descriptive data for the personal, anthropometric, biomechanical and training-related characteristics
292 will be presented as count and percentage for dichotomous variables, and as mean and standard
293 deviation, or as median and range, respectively, for normally and non-normally distributed continuous
294 variables. Average sport-related characteristics will be computed for each participant over their

specific period of observation. Shock absorption properties (stiffness, N/mm) of the two types of shoes will be compared using a Student's t test.

Cox proportional hazards regressions will be used to compute the hazard rates in the exposure groups, using first-time injury as the primary outcome. Date of inclusion (baseline evaluation date) and date of injury or of censoring will be basic data used to calculate the time at risk, which is expressed in hours spent running and defined as the time-scale.[35] A participant will be right-censored if injury unrelated to running or severe disease caused a modification of the running plan, or at the end of follow-up. Reasons for right-censoring will be reported. The assumption of proportional hazards will be evaluated by log-minus-log plots.

Unadjusted Cox regressions will be performed to present the crude estimates of HRs for shoe model, body mass and other potential risk factors such as running biomechanics variables (see table 1) and training-related characteristics. Body mass is an exposure that can change over time (time-dependent covariate). This means that each participant could move between exposure states continuously (every month in our study). A delayed entry will be used in the unadjusted Cox regression model for body mass.[46]

Subsequently, the variables with a P value <0.200 will be included in the adjusted Cox regression analysis to determine whether shoe cushioning and/or body mass are associated with injury risk, controlling for potential confounders. The recommendation for using at least 10 injuries per predictor variable included in the Cox regression analysis will be strictly followed.[47]

Finally, to investigate if the effect of shoe cushioning on RRI risk is modified by body mass, a stratified analysis will be performed using the median value of body mass as cut-off. HRs and their 95% confidence intervals (CI) will be determined within each stratum.[48] All analyses will be performed using STATA/SE version 14.

DISCUSSION

It is common belief that shoe cushioning technology protects the runner against harmful consequences of repetitive high-load impacts. Therefore, heavier runners are generally advised to use footwear with

adapted shock absorption properties. Surprisingly, few studies have investigated the impact of shoe cushioning on injury risk.[32, 33] These studies did not provide any evidence on the beneficial effect of increased shock absorption properties. However, none of them included anthropometric measures in their analyses. Also, one study compared different types of insoles added in the shoes,[33] while the other compared two versions of a standard running shoe with a limited difference in midsole hardness (~15%).[32] Other study limitations such as the sample size ($n < 250$)[32] or the study population (Royal Air Force recruits)[33] suggest that these results should be interpreted with caution. The evidence on the association between running shoe cushioning and RRI is still poor and inconclusive. One of the main reasons is the practical constraint of investigations trying to combine biomechanical analyses with a long-term prospective follow-up in a large number of runners.[11] This study is the first randomised controlled trial investigating the influence of shoe cushioning on RRI risk including an evaluation of running technique in all participants. The results will provide information on the real benefits provided by additional cushioning, as well as on the mechanisms that might explain any potential preventive effect.

ETHICS AND DISSEMINATION

This study will be conducted in accordance with the Declaration of Helsinki and the Medical Research Involving Human Subjects Act. Also, the study protocol (Ref: 201701/02 v1.1) was approved by the National Ethics Committee for Research (www.cner.lu). Written informed consent will be obtained from all participants (Supplementary file 4). All collected data will be stored electronically using a coding system. This will ensure that the data is used in the strictest confidence and will not reveal the identity of the participants. Collected raw data will not be passed on to unauthorised third parties. Results presented or published in articles and reports will be depicted in general terms, to maintain participant anonymity. Electronic data will be stored on a secure server in data files only accessible to the project leader and co-investigators of the project. A notification of this study was sent to the National Data Protection Agency (CNPD). Study results will be submitted for

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

publication in peer-reviewed journals and for presentation at international conferences. Furthermore,
we aim to disseminate our results through popular specialised magazines and websites.

Contributors - LM, ND, AU and DT contributed to the study conception and study design. LM is the
main investigator, wrote the article with input from other investigators, and will be responsible for the
acquisition and analysis of the data. ND will be responsible for the shoe design, production and
testing. LM and DT will be responsible for data interpretation and manuscript drafting. ND, AU and
DT commented on the various versions of the study protocol. All authors approved the final
manuscript.

Funding – This study is co-funded by Decathlon, Villeneuve d’Ascq, France and by the Ministry of
Higher Education and Research of Luxembourg.

Competing interests – A research partnership agreement was signed between Decathlon and the
Luxembourg Institute of Health (LIH). ND is employed at Decathlon SA. Decathlon will not be
involved in the collection, management, analysis and interpretation of data. LM, DT and AU may not
gain or lose financially from the results of the study in any way.

Ethics approval - All procedures were approved by the National Ethics Committee for Research
(Ref: 201701/02 v1.1).

REFERENCE LIST

- 1 Kohl HW, 3rd, Craig CL, Lambert EV, et al. The pandemic of physical inactivity: global action for public health. *Lancet*. 2012;**380**:294-305.
- 2 World Health Organization (WHO). Global status report on noncommunicable diseases 2014. . *Geneva: World Health Organization (WHO)*. 2014.
- 3 Hespanhol Junior LC, Pillay JD, van Mechelen W, et al. Meta-Analyses of the Effects of Habitual Running on Indices of Health in Physically Inactive Adults. *Sports Med*. 2015;**45**:1455-68.
- 4 Lopes AD, Hespanhol Junior LC, Yeung SS, et al. What are the main running-related musculoskeletal injuries? A Systematic Review. *Sports Med*. 2012;**42**:891-905.
- 5 Alentorn-Geli E, Samuelsson K, Musahl V, et al. The Association of Recreational and Competitive Running With Hip and Knee Osteoarthritis: A Systematic Review and Meta-Analysis. *J Orthop Sports Phys Ther*. 2017:1-36.
- 6 Koplan JP, Rothenberg RB, Jones EL. The natural history of exercise: a 10-yr follow-up of a cohort of runners. *Med Sci Sports Exerc*. 1995;**27**:1180-4.
- 7 Hespanhol Junior LC, van Mechelen W, Postuma E, et al. Health and economic burden of running-related injuries in runners training for an event: A prospective cohort study. *Scand J Med Sci Sports*. 2016;**26**:1091-9.
- 8 Hespanhol Junior LC, van Mechelen W, Verhagen E. Health and Economic Burden of Running-Related Injuries in Dutch Trailrunners: A Prospective Cohort Study. *Sports Med*. 2016.
- 9 van Gent RN, Siem D, van Middelkoop M, et al. Incidence and determinants of lower extremity running injuries in long distance runners: a systematic review. *Br J Sports Med*. 2007;**41**:469-80; discussion 80.

1
2
3 393 10 Knapik JJ, Jones BH, Steelman RA. Physical training in boots and running shoes: a
4
5 394 historical comparison of injury incidence in basic combat training. *Mil Med.* 2015;**180**:321-8.
6
7 395 11 Theisen D, Malisoux L, Gette P, et al. Footwear and running-related injuries –Running on
8
9 396 faith? *Sports Orthop Traumatol.* 2016;**32**:169-76.
10
11 397 12 Knapik JJ, Trone DW, Tchandja J, et al. Injury-reduction effectiveness of prescribing
12
13 398 running shoes on the basis of foot arch height: summary of military investigations. *J Orthop*
14
15 399 *Sports Phys Ther.* 2014;**44**:805-12.
16
17 400 13 Meeuwisse WH, Tyreman H, Hagel B, et al. A dynamic model of etiology in sport injury:
18
19 401 the recursive nature of risk and causation. *Clin J Sport Med.* 2007;**17**:215-9.
20
21 402 14 Hreljac A. Impact and overuse injuries in runners. *Med Sci Sports Exerc.* 2004;**36**:845-9.
22
23 403 15 Malisoux L, Nielsen RO, Urhausen A, et al. A step towards understanding the
24
25 404 mechanisms of running-related injuries. *J Sci Med Sport.* 2015;**18**:523-8.
26
27 405 16 van der Worp H, Vrielink JW, Bredeweg SW. Do runners who suffer injuries have higher
28
29 406 vertical ground reaction forces than those who remain injury-free? A systematic review and
30
31 407 meta-analysis. *Br J Sports Med.* 2016.
32
33 408 17 Davis IS, Bowser BJ, Mullineaux DR. Greater vertical impact loading in female runners
34
35 409 with medically diagnosed injuries: a prospective investigation. *Br J Sports Med.* 2015.
36
37 410 18 Barton CJ, Bonanno DR, Carr J, et al. Running retraining to treat lower limb injuries: a
38
39 411 mixed-methods study of current evidence synthesised with expert opinion. *Br J Sports Med.*
40
41 412 2016;**50**:513-26.
42
43 413 19 Diebal AR, Gregory R, Alitz C, et al. Forefoot running improves pain and disability
44
45 414 associated with chronic exertional compartment syndrome. *Am J Sports Med.* 2012;**40**:1060-
46
47 415 7.
48
49
50
51
52
53
54
55
56
57
58
59
60

- 20 Azevedo AP, Mezencio B, Amadio AC, et al. 16 Weeks of Progressive Barefoot Running Training Changes Impact Force and Muscle Activation in Habitual Shod Runners. *PLoS One*. 2016;**11**:e0167234.
- 21 Boyer ER, Derrick TR. Select injury-related variables are affected by stride length and foot strike style during running. *Am J Sports Med*. 2015;**43**:2310-7.
- 22 Luedke LE, Heiderscheit BC, Williams DS, et al. Influence of Step Rate on Shin Injury and Anterior Knee Pain in High School Runners. *Med Sci Sports Exerc*. 2016;**48**:1244-50.
- 23 Lorimer AV, Hume PA. Stiffness as a Risk Factor for Achilles Tendon Injury in Running Athletes. *Sports Med*. 2016;**46**:1921-38.
- 24 Baltich J, Maurer C, Nigg BM. Increased vertical impact forces and altered running mechanics with softer midsole shoes. *PLoS One*. 2015;**10**:e0125196.
- 25 Hardin EC, van den Bogert AJ, Hamill J. Kinematic adaptations during running: effects of footwear, surface, and duration. *Med Sci Sports Exerc*. 2004;**36**:838-44.
- 26 Nigg BM, Baltich J, Maurer C, et al. Shoe midsole hardness, sex and age effects on lower extremity kinematics during running. *J Biomech*. 2012;**45**:1692-7.
- 27 Shorten M, Mientjes M. The 'heel impact' force peak during running is neither 'heel' nor 'impact' and does not quantify shoe cushioning effects. *Footwear Science*. 2011;**3**:41-58.
- 28 Logan S, Hunter I, JT JTH, et al. Ground reaction force differences between running shoes, racing flats, and distance spikes in runners. *J Sports Sci Med*. 2010;**9**:147-53.
- 29 O'Leary K, Vorpahl KA, Heiderscheit B. Effect of cushioned insoles on impact forces during running. *J Am Podiatr Med Assoc*. 2008;**98**:36-41.
- 30 Kersting UG, Bruggemann GP. Midsole material-related force control during heel-toe running. *Res Sports Med*. 2006;**14**:1-17.
- 31 Nigg BM, Bahlsten HA, Luethi SM, et al. The influence of running velocity and midsole hardness on external impact forces in heel-toe running. *J Biomech*. 1987;**20**:951-9.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

441 32 Theisen D, Malisoux L, Genin J, et al. Influence of midsole hardness of standard
442 cushioned shoes on running-related injury risk. *Br J Sports Med.* 2014;**48**:371-6.

443 33 Withnall R, Eastaugh J, Freemantle N. Do shock absorbing insoles in recruits undertaking
444 high levels of physical activity reduce lower limb injury? A randomized controlled trial. *J R*
445 *Soc Med.* 2006;**99**:32-7.

446 34 Buist I, Bredeweg SW, Bessem B, et al. Incidence and risk factors of running-related
447 injuries during preparation for a 4-mile recreational running event. *Br J Sports Med.*
448 2010;**44**:598-604.

449 35 Nielsen RO, Malisoux L, Moller M, et al. Shedding Light on the Etiology of Sports
450 Injuries: A Look Behind the Scenes of Time-to-Event Analyses. *J Orthop Sports Phys Ther.*
451 2016:1-34.

452 36 Malisoux L, Chambon N, Delattre N, et al. Injury risk in runners using standard or motion
453 control shoes: a randomised controlled trial with participant and assessor blinding. *Br J*
454 *Sports Med.* 2016;**50**:481-7.

455 37 Malisoux L, Chambon N, Urhausen A, et al. Influence of the Heel-to-Toe Drop of
456 Standard Cushioned Running Shoes on Injury Risk in Leisure-Time Runners: A Randomized
457 Controlled Trial With 6-Month Follow-up. *Am J Sports Med.* 2016.

458 38 International A. ASTM F1614-99 Standard test method for shock attenuating properties of
459 materials systems for athletic footwear. *ASTM International.* 2006:West Conshohocken, PA,
460 USA.

461 39 Delattre N, Chambon N, Berton E, et al. Effect of time during a running session with
462 minimal footwear. *Comput Methods Biomech Biomed Engin.* 2013;**16 Suppl 1**:104-5.

463 40 Sabharwal S, Kumar A. Methods for assessing leg length discrepancy. *Clin Orthop Relat*
464 *Res.* 2008;**466**:2910-22.

- 465 41 Morin JB, Dalleau G, Kyrolainen H, et al. A simple method for measuring stiffness during
466 running. *J Appl Biomech*. 2005;**21**:167-80.
- 467 42 Malisoux L, Ramesh J, Mann R, et al. Can parallel use of different running shoes decrease
468 running-related injury risk? *Scand J Med Sci Sports*. 2015;**25**:110-5.
- 469 43 Foster C. Monitoring training in athletes with reference to overtraining syndrome. *Med Sci*
470 *Sports Exerc*. 1998;**30**:1164-8.
- 471 44 Yamato TP, Saragiotto BT, Lopes AD. A consensus definition of running-related injury in
472 recreational runners: a modified Delphi approach. *J Orthop Sports Phys Ther*. 2015;**45**:375-
473 80.
- 474 45 Rae K, Orchard J. The Orchard Sports Injury Classification System (OSICS) version 10.
475 *Clin J Sport Med*. 2007;**17**:201-4.
- 476 46 Nielsen RO, Parner ET, Nohr EA, et al. Excessive progression in weekly running distance
477 and risk of running-related injuries: an association which varies according to type of injury. *J*
478 *Orthop Sports Phys Ther*. 2014;**44**:739-47.
- 479 47 Peduzzi P, Concato J, Feinstein AR, et al. Importance of events per independent variable
480 in proportional hazards regression analysis. II. Accuracy and precision of regression
481 estimates. *J Clin Epidemiol*. 1995;**48**:1503-10.
- 482 48 Knol MJ, VanderWeele TJ. Recommendations for presenting analyses of effect
483 modification and interaction. *Int J Epidemiol*. 2012;**41**:514-20.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


FIGURE LEGEND

Figure 1: Trial design.

SUPPLEMENTARY FILES

- Supplementary file 1: SPIRIT Checklist
- Supplementary file 1: Study schedule
- Supplementary file 3: Risk of sport participation form
- Supplementary file 4: Informed consent

Figure 1: Trial Design



Trial design



SPIRIT 2013 Checklist: Recommended items to address in a clinical trial protocol and related documents*

Section/item	Item No	Description	Addressed on page number
Administrative information			
Title	1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	Page 1
Trial registration	2a	Trial identifier and registry name. If not yet registered, name of intended registry	Pages 2 & 7: clinicaltrials.gov (NCT03115437)
	2b	All items from the World Health Organization Trial Registration Data Set	/
Protocol version	3	Date and version identifier	Page 7
Funding	4	Sources and types of financial, material, and other support	Page 15
Roles and responsibilities	5a	Names, affiliations, and roles of protocol contributors	Page 15
	5b	Name and contact information for the trial sponsor	Page 1 & 15
	5c	Role of study sponsor and funders, if any, in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication, including whether they will have ultimate authority over any of these activities	Page 15

5d	Composition, roles, and responsibilities of the coordinating centre, steering committee, endpoint adjudication committee, data management team, and other individuals or groups overseeing the trial, if applicable (see Item 21a for data monitoring committee)	n.a.
----	--	------

Introduction

Background and rationale	6a	Description of research question and justification for undertaking the trial, including summary of relevant studies (published and unpublished) examining benefits and harms for each intervention	Pages 4 and 5
	6b	Explanation for choice of comparators	Pages 5 and 4
Objectives	7	Specific objectives or hypotheses	Page 6
Trial design	8	Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group), allocation ratio, and framework (eg, superiority, equivalence, noninferiority, exploratory)	Page 7 and 8

Methods: Participants, interventions, and outcomes

Study setting	9	Description of study settings (eg, community clinic, academic hospital) and list of countries where data will be collected. Reference to where list of study sites can be obtained	Page 7
Eligibility criteria	10	Inclusion and exclusion criteria for participants. If applicable, eligibility criteria for study centres and individuals who will perform the interventions (eg, surgeons, psychotherapists)	Page 7
Interventions	11a	Interventions for each group with sufficient detail to allow replication, including how and when they will be administered	Page 8
	11b	Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose change in response to harms, participant request, or improving/worsening disease)	Page 11-12
	11c	Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence (eg, drug tablet return, laboratory tests)	Page 11-13
	11d	Relevant concomitant care and interventions that are permitted or prohibited during the trial	n.a.

1	Outcomes	12	Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg, median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen efficacy and harm outcomes is strongly recommended	Page 9 and 11
2				
3				
4				
5				
6	Participant timeline	13	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for participants. A schematic diagram is highly recommended (see Figure)	Figure 1 and Suppl. file 2
7				
8				
9				
10	Sample size	14	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	Page 12
11				
12				
13	Recruitment	15	Strategies for achieving adequate participant enrolment to reach target sample size	Page 7
14				
15				

16 **Methods: Assignment of interventions (for controlled trials)**

17 Allocation:

18				
19				
20	Sequence generation	16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	Page 8
21				
22				
23				
24				
25				
26	Allocation concealment mechanism	16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	Page 8
27				
28				
29				
30	Implementation	16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	Page 8
31				
32				
33				
34	Blinding (masking)	17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	Page 8
35				
36				
37		17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	Page 8
38				
39				
40				

41 **Methods: Data collection, management, and analysis**

1	Data collection	18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related	Pages 9 to 11
2	methods		processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of	
3			study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known.	
4			Reference to where data collection forms can be found, if not in the protocol	
5				
6		18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be	Page 11
7			collected for participants who discontinue or deviate from intervention protocols	
8				
9	Data management	19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality	Page 12
10			(eg, double data entry; range checks for data values). Reference to where details of data management	
11			procedures can be found, if not in the protocol	
12				
13	Statistical methods	20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the	Pages 12 and 13
14			statistical analysis plan can be found, if not in the protocol	
15				
16		20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	Pages 12 and 13
17				
18		20c	Definition of analysis population relating to protocol non-adherence (eg, as randomised analysis), and any	Pages 12 and 13
19			statistical methods to handle missing data (eg, multiple imputation)	
20				
21				
22				
23				
24	Methods: Monitoring			
25				
26	Data monitoring	21a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of /	
27			whether it is independent from the sponsor and competing interests; and reference to where further details	
28			about its charter can be found, if not in the protocol. Alternatively, an explanation of why a DMC is not	
29			needed	
30				
31		21b	Description of any interim analyses and stopping guidelines, including who will have access to these interim /	
32			results and make the final decision to terminate the trial	
33				
34				
35	Harms	22	Plans for collecting, assessing, reporting, and managing solicited and spontaneously reported adverse	Page 12
36			events and other unintended effects of trial interventions or trial conduct	
37				
38	Auditing	23	Frequency and procedures for auditing trial conduct, if any, and whether the process will be independent /	
39			from investigators and the sponsor	
40				
41				

Ethics and dissemination

1	Research ethics	24	Plans for seeking research ethics committee/institutional review board (REC/IRB) approval	Page 14
2	approval			
3				
4	Protocol	25	Plans for communicating important protocol modifications (eg, changes to eligibility criteria, outcomes,	Page 14
5	amendments		analyses) to relevant parties (eg, investigators, REC/IRBs, trial participants, trial registries, journals,	
6			regulators)	
7				
8	Consent or assent	26a	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and	Page 7
9			how (see Item 32)	
10				
11				
12		26b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary	n.a.
13			studies, if applicable	
14				
15	Confidentiality	27	How personal information about potential and enrolled participants will be collected, shared, and maintained	Page 7
16			in order to protect confidentiality before, during, and after the trial	
17				
18	Declaration of	28	Financial and other competing interests for principal investigators for the overall trial and each study site	Page 15
19	interests			
20				
21	Access to data	29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that	Pages 14
22			limit such access for investigators	
23				
24				
25	Ancillary and post-	30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial	/
26	trial care		participation	
27				
28	Dissemination policy	31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals,	Page 14
29			the public, and other relevant groups (eg, via publication, reporting in results databases, or other data	
30			sharing arrangements), including any publication restrictions	
31				
32				
33		31b	Authorship eligibility guidelines and any intended use of professional writers	/
34				
35		31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	n.a.
36				
37				
38	Appendices			
39				
40	Informed consent	32	Model consent form and other related documentation given to participants and authorised surrogates	Supplementary file
41	materials			4
42				
43				
44				
45				
46				
47				

1 Biological 33 Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular n.a.
2 specimens analysis in the current trial and for future use in ancillary studies, if applicable
3

4 *It is strongly recommended that this checklist be read in conjunction with the SPIRIT 2013 Explanation & Elaboration for important clarification on the items.
5 Amendments to the protocol should be tracked and dated. The SPIRIT checklist is copyrighted by the SPIRIT Group under the Creative Commons
6 [“Attribution-NonCommercial-NoDerivs 3.0 Unported”](#) license.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

For peer review only

Supplementary file 2: Schedule of enrolment, interventions, and assessments for the study

	STUDY PERIOD							
	Enrolment	Allocation	Post-allocation: For every sport activity					Close-out
TIMEPOINT	-t ₁	0	t ₁	t ₂	t ₃	t ₄	etc.	6 Months
ENROLMENT								
Eligibility screen	X							
Informed consent	X							
Baseline Questionnaire	X	X						
Allocation		X						
Shoe distribution		X						
Running analysis		X						
INTERVENTIONS								
[Intervention A]								
[Intervention B]								
ASSESSMENTS								
Running experience	X	X						
Running regularity	X	X						
Typical running frequency	X	X						
Typical running distance	X	X						
Training running speed	X	X						
Type of running	X	X						
Competition participation	X	X						
Last event distance	X	X						
Favourite running distance	X	X						
Best time on 5 km / 10km	X	X						
Previous injury	X	X						

Height	X	X						
Body mass	X	X						
Leg length	X	X						
% fat tissue	X	X						
Step frequency	X	X						
Contact time	X	X						
Flight time	X	X						
Duty factor	X	X						
Step length	X	X						
Vertical Impact Peak Force	X	X						
Peak Vertical Force	X	X						
Vertical Instantaneous Loading Rate	X	X						
Vertical Average Loading Rate	X	X						
Peak Power	X	X						
Time to Peak Force	X	X						
Leg stiffness	X	X						
Vertical stiffness	X	X						
Sports discipline			X	X	X	X	etc.	
Duration			X	X	X	X	etc.	
Distance (if applicable)			X	X	X	X	etc.	
Perceived Exertion			X	X	X	X	etc.	
Shoe used (if running)			X	X	X	X	etc.	
Surface (if running)			X	X	X	X	etc.	
Pain*			X	X	X	X	etc.	
Injury**			X	X	X	X	etc.	

*The pain did not stop the participant from continuing normal training

**The participants had to adapt or interrupt their training accordingly

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplementary file 4 : Questionnaire on risk of sport participation

- 1. Past medical history, have you had:
 - 1.1 Severe cardiac arrhythmia?
 - 1.2 Myocardial infarction?
 - 1.3 Heart surgery?
 - 1.4 Intracardiac catheter?
 - 1.5 Coronary angioplasty (dilatation by balloon, stenting)?
 - 1.6 Pacemaker or heart defibrillator?
 - 1.7 Cardiac insufficiency?
 - 1.8 Heart transplantation?
 - 1.9 Congenital heart defect?
- 2. Past and present complaints
 - 2.1 Chest pain / discomfort during physical exertion?
 - 2.2 Dyspnea (uncommon breathlessness)?
 - 2.3 Dizziness/unconsciousness?
 - 2.4 Palpitations, tachycardia, pulse irregularities?
 - 2.5 Intake of any cardiac drugs?
- 3. Other disorders
 - 3.1 Muscular or articular complaints?
 - 3.2 Other drugs?
 - 3.3 Insecurity during physical exertion?
 - 3.4 For females: Pregnancy?
- 4. Cardiovascular risk factors
 - 4.1 Are you male over 45 years?
 - 4.2 Are you female over 55 years or you have had a hysterectomy or you are menopausal?
 - 4.3 Smoker (active / in the past 10 years)?
 - 4.4 Your blood pressure is over 140/90 mmHg or you take antihypertensive drugs?
 - 4.5 Your cholesterol level is over 240 mg/dl?
 - 4.6 Myocardial infarction, stroke, marfan disease or sudden cardiac death in the family ? (Father resp. brother before age of 55 years/ Mother resp. sister before 65 years)?
 - 4.7 You are diabetic or you take antidiabetic drugs?
 - 4.8 Sports activity less than 90 min/week?
 - 4.9 You have a Body Mass Index (BMI) over 30 ?

Free Informed Consent

Title:

Institution:

Project manager:

Research assistant:

Head of unit:

1. I declare to have read the above-described information and accept to voluntarily participate in the study "Effects of bodyweight and shoe cushioning on injury risk and running biomechanics: A randomised control trial" conducted by the SMRL.
2. I accept that my data shall be used and communicated to the commercial partner for strictly scientific purposes once it has been pseudonymised (coded).
3. I received a copy of the present signed informed consent document, as well as the general information intended for athlete participants. I received a clear description of the purpose and the nature of the study and I am aware of what is expected of me as a participant in this study. I have had enough time and the opportunity to ask questions about the study; all my questions have been met with a satisfactory answer.
4. I am free to retire from the study at any time without justification. By doing so I will not suffer any material or moral damage.
5. I agree that the results of this study can be subject to public talks or scientific publication.
6. I voluntarily consent to participate in this study and I fully understand what kind of data will be gathered during the study.
7. I preserve/abide the rights of access, deletion or modification of my personal data. Any personal information will be kept confidential and protected in agreement with the modified personal data protection act of August 2nd 2002. I can exercise that right via the project manager.

The responding signatory freely consents to participate in the above mentioned study

Name and First Name of the respondent:

Signature of the respondent:

Name and signature of the project manager:

Place and date:

BMJ Open

Shoe cushioning, body mass and running biomechanics as risk factors for running injury: a study protocol for a randomised controlled trial

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-017379.R2
Article Type:	Protocol
Date Submitted by the Author:	04-Jul-2017
Complete List of Authors:	Malisoux, Laurent; Luxembourg Institute of Health, Department of Population Health Delattre, Nicolas; Decathlon SportsLab, Movement Sciences Department Urhausen, Axel; Luxembourg Institute of Health, Department of Population Health; Centre Hospitalier de Luxembourg, Sports Clinic Theisen, Daniel; Luxembourg Institute of Health, Department of Population Health
Primary Subject Heading:	Sports and exercise medicine
Secondary Subject Heading:	Public health
Keywords:	Sports injury prevention, Footwear, Impact forces, EPIDEMIOLOGY

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 TITLE PAGE

2 Title:

3 Shoe cushioning, body mass and running biomechanics as risk factors for running injury: a study
4 protocol for a randomised controlled trial

5 Authors:

6 Laurent MALISOUX^a, Nicolas DELATTRE^b, Axel URHAUSEN^{a,c,d}, Daniel THEISEN^a

7 Affiliations:

8 ^a Luxembourg Institute of Health, Sports Medicine Research Laboratory, Luxembourg, Grand-Duchy
9 of Luxembourg

10 ^b Decathlon SportsLab, Movement Sciences Department, Villeneuve d’Ascq, France

11 ^c Centre Hospitalier de Luxembourg, Sports Clinic, Luxembourg, Grand-Duchy of Luxembourg

12 ^d International University of Health, Exercise and Sports (LUNEX), 50 avenue du Parc des Sports L-
13 4671 Differdange, Grand-Duchy of Luxembourg

14 Corresponding Author:

15 Laurent Malisoux

16 76 rue d’Eich

17 L-1460 Luxembourg

18 E-Mail: laurent.malisoux@lih.lu

19 Phone: +352 26970 231

20 Keywords:

21 Sports injury prevention, footwear, epidemiology, impact forces

22 Word count (excluding title page, abstract, references, figures and tables): 3476

23 Abstract Word count: 260

24 Number of figures: 1

25 Number of tables: 1

26 Online supplementary material: 2

27

Title: Shoe cushioning, body mass and running biomechanics as risk factors for running injury: a study protocol for a randomised controlled trial

ABSTRACT

Introduction: Repetitive loading of the musculoskeletal system is suggested to be involved in the underlying mechanism of the majority of running-related injuries (RRI). Accordingly, heavier runners are assumed to be at a higher risk of RRI. The cushioning system of modern running shoes is expected to protect runners against high impact forces, and therefore, RRI. However, the role of shoe cushioning in injury prevention remains unclear. The main aim of this study is to investigate the influence of shoe cushioning and body mass on RRI risk, while exploring simultaneously the association between running technique and RRI risk.

Methods and analysis: This double-blinded randomised controlled trial will involve about 800 healthy leisure-time runners. They will randomly receive one of two running shoe models that will differ in their cushioning properties (i.e. stiffness) by ~35%. The participants will perform a running test on an instrumented treadmill at their preferred running speed at baseline. They will then be followed-up prospectively over a 6-month period, during which they will self-report all their sports activities as well as any injury in an internet-based database TIPPS (Training and Injury Prevention Platform for Sports). Cox regression analyses will be used to compare injury risk between the study groups and to investigate the association between training, biomechanical and anatomical risk factors, and injury risk.

Ethics and dissemination: The study was approved by the National Ethics Committee for Research (Ref: 201701/02 v1.1). Outcomes will be disseminated through publications in peer-reviewed journals, presentations at international conferences, as well as articles in popular magazines and on specialised websites.

Trial registration number: NCT03115437

1
2
3 55
4
5 56
6
7 57
8
9 58
10
11 59
12
13 60
14
15 61
16
17 62
18
19 63
20
21 64
22
23 65
24
25 66
26
27 67
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STRENGTHS AND LIMITATIONS OF THIS STUDY

- Double-blinded randomised controlled trial (assessor and participant blinding) and intention-to-treat analysis.
- This study compares 2 shoe versions with widely differing cushioning properties while remaining within the cushioning range of models available on the market.
- A biomechanical analysis will be performed for each participant prior to the 6-month follow-up, which allows to investigate the association between running biomechanics and injury risk in a large cohort of runners.
- The running test will be carried out on a treadmill using a standardised protocol, which might not be reflective of the participants' habitual training conditions.

INTRODUCTION

Running is an increasingly popular form of physical activity. From a public health perspective, the promotion of leisure-time running might be a powerful strategy to combat the pandemic of physical inactivity worldwide,[1] and its consequence on non-communicable diseases.[2] Although regular running activity has a massive beneficial impact on health,[3] it also generates a relatively high number of injuries, especially at the lower limb.[4] The risk of sustaining a running-related injury (RRI) cancels out part of the benefits of running practice, since the long term consequences of injury might include, among others, increased risk of osteoarthritis,[5] a reduction of physical activity,[6] as well as an increase in health care costs.[7, 8] RRI incidence has remained high during the last 40 years, with an overall incidence rate ranging between 18.2% and 92.4%.[9] The role of footwear on RRI risk has been strongly emphasized ever since jogging became popular in the 1970s, but there is currently no evidence that developments in running shoe technology and new concepts regularly emerging on the market have helped to tackle the RRI burden.[10-12]

Most RRI are overuse injuries, as they develop progressively over the kilometres run. The aetiology of these injuries is multifactorial,[13] which implies that to understand the mechanisms leading to injury, a holistic approach is warranted, including the study of a large set of potential risk factors. These factors could be classified as being related to training characteristics, running mechanics and anatomy of the runners. Some authors suggested that anatomical and biomechanical factors influence the tolerance to physical strain and thus the relationship between training load and injury occurrence.[14, 15]

Biological tissues such as bones, muscles and tendons can endure a certain amount of stress, provided that the product of stress level (e.g. intensity, external load) and the number of repetitions within a certain time period (e.g. strides, training sessions) remains below a threshold that is specific to each structure.[14] In running, the ground reaction force is the main external stress that acts on the body. Vertical ground reaction force (VGRF) is a biomechanical factor that has been extensively studied in

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

95 running.[16, 17] A recent meta-analysis found that the loading rate of the vertical ground reaction
96 force was higher in patients with a history of stress fracture.[16] High impact-related variables were
97 shown to increase the risk of bony and soft tissue injuries.[17] Moreover, running retraining
98 interventions have proven their efficiency in modifying some VGRF parameters and decreasing pain,
99 which suggest that running retraining represents an interesting paradigm to treat RRI.[18-20] Other
100 biomechanical factors such as step length,[21] step frequency [22] or leg stiffness [23] have
101 previously been suggested as potential biomechanical risk factors for RRI, yet no causal relationship
102 has been established.
103
104 Since running biomechanics are associated with injury risk, any effect of shoe features on the running
105 pattern and VGRF parameters deserve attention. Given that repetitive loading of the musculoskeletal
106 system is an injury risk factor, cushioning has been one of the most extensively investigated shoe
107 features. The shock absorption properties of footwear mainly result from the materials used in the sole
108 (i.e. their type, density, structure and combination), as well as from the geometry of the shoe (i.e. the
109 midsole thickness and the design of inserts). One of the most popular approaches has been to change
110 the hardness of the shoe midsole.[24-26] Overall, the studies investigating the effect of shoe
111 cushioning on VGRF did not provide consistent results. In theory, peak impact forces should be
112 reduced by softer or more compliant shoes,[27] which was indeed confirmed in some in vivo
113 studies.[28, 29] Conversely, some investigations did not find any effect of cushioning,[30] or reported
114 increased peak impact forces in softer shoes.[24, 31] Recently, a large cross sectional study revealed
115 that softer midsole hardness was associated with higher vertical force impact peak.[24] Unfortunately,
116 very few studies have investigated the association between shoe cushioning and injury risk.[32, 33] In
117 a previous randomised controlled trial, midsole hardness was not associated with RRI risk. However,
118 the difference in shoe stiffness between the shoe conditions was limited (15%).[32] Therefore, the
119 role of shoe cushioning systems in RRI prevention remains unclear.

1
2
3 121 Body mass index (BMI) has been associated with injury risk in novice,[34, 35] as well as in
4
5 122 recreational runners,[32] though other results suggest a protective effect of BMI.[9] It is common
6
7 123 belief that individuals with higher BMI have a higher injury risk, because of the increased physical
8
9 124 stress that results from extra body weight. Surprisingly, body mass as such has hardly ever been
10
11 125 considered as a potential risk factor for running injury.[9]
12

13
14 126

15 127 Surprisingly, the literature on the association between single shoe features and RRI risk is still
16
17 128 poor.[11, 36, 37] Until now, no relationship has been found between the cushioning properties of
18
19 129 modern running shoes and RRI risk,[32] but body mass should be taken into account here. Therefore,
20
21 130 the main purpose of this study is to investigate the association between shoe cushioning and body
22
23 131 mass on the one hand, and RRI risk on the other hand in recreational runners. The secondary aims are
24
25 132 to identify which of the running technique-related characteristics (timing variables and VGRF
26
27 133 parameters) are associated with injury risk, as well as with the cushioning properties of the shoes.
28
29 134 Shoe cushioning will be characterised by the stiffness at the heel (N/mm) and quantified by
30
31 135 standardised impact test.[38] The following hypotheses (H) will be tested:

32
33 136 H1. Running shoes with greater stiffness are associated with a higher injury risk in leisure-time
34
35 137 runners.

36
37 138 H2. High body mass is associated with a higher injury risk in leisure-time runners.

38
39 139 H3. Runners with a high body mass experience a lower injury risk in shoes with greater stiffness.

40
41 140 H4. A higher step length, a lower step frequency, and higher vertical loading rate are associated with a
42
43 141 higher injury risk.

44
45 142 H5. Running shoes with greater stiffness will be associated with higher vertical loading rate and a
46
47 143 shorter contact time.

48
49 144 H6. High body mass will be associated with higher vertical loading rate, increased contact time,
50
51 145 increased duty factor, and decreased step frequency.

52
53 146 Furthermore, exploratory risk factor analyses will be performed on the biomechanical variables
54
55 147 obtained from the running analysis, anthropometric measurements, running experience, and habitual
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

running speed. The focus of the analyses is the effect modification of body mass and other above mentioned risk factors on the association between shoe cushioning and injury risk.

METHODS AND ANALYSIS

Trial design

The design of this study is a randomised controlled trial with a 6-month follow-up and a biomechanical analysis of running pattern at baseline. The study is based on the comparison between 2 running shoe prototypes, which only differ with respect to the cushioning (i.e. stiffness). The cushioning properties of both shoe versions are within the range of those from available models on the market. Running footwear is provided by a renowned sport equipment manufacturer. The main outcome is RRI (cf. definition below). The participants as well as the assessors are blinded to group allocation. The design of the trial is illustrated in Figure 1. The protocol conforms to the Recommendations for Interventional Trials (SPIRIT) and has been registered on <https://clinicaltrials.gov/> (NCT03115437, 11/04/2017).

Insert Figure 1 about here

Study population

The target population is leisure-time runners, regardless of running experience, fitness level, or body mass. Participants will be recruited through advertisements in local newspapers, social media, running magazines and press releases within the country during the months of September 2017 to January 2018. Healthy volunteers will be considered eligible if they are aged between 18 and 65 years and capable of performing 15 minutes of consecutive running. Volunteers will be excluded in case of any contraindication to perform running activity, prior (<12 months) surgery or major trauma to the lower limbs or lower back region, any running impeding injury over the previous month, or use of orthopaedic insoles for running activities. Additionally, the participants will have to agree on the following requirements: 1) to practice running at least once a week, 2) to use the provided study shoes

1
2
3 175 for all their running sessions, and 3) to report, at least once per week, all sports activities, as well as
4
5 176 any injury or pain experienced during the follow-up period on an internet-based database called
6
7 177 TIPPS (Training and Injury Prevention Platform for Sports, www.tipps.lu). Volunteers first have to
8
9 178 create a personal account on TIPPS, pre-register to the study via their personal account, and answer an
10
11 179 online inclusion/exclusion questionnaire as well as a baseline questionnaire. Answers to both
12
13 180 questionnaires will be assessed by the investigators during the initial visit.
14
15
16

181

182 **Randomisation**

183 Participants must understand and agree on the randomized design of the study. Those who meet the
184 eligibility criteria and sign the informed consent form will be randomly allocated to one of the two
185 study arms. They will be stratified according to their sex, which is known to influence body mass as
186 well as many other anthropometric characteristics. Therefore, two pre-established randomisation lists
187 (block size = 40) will be prepared by a statistician not involved in any other part of the study before
188 the beginning of the recruitment. To ensure allocation concealment, the study groups and shoes will
189 be coded and the randomisation lists will be uploaded in the TIPPS system by an IT specialist who
190 will not be involved in any other part of the study. Then, the TIPPS system will provide the
191 investigator in charge of the recruitment with a study group number for each participant, according to
192 the randomisation lists. The investigator will upload the shoe number according to shoe size chosen
193 and study arm so that a cross validation will be performed by the electronic system. The investigators
194 in charge of the recruitment, the follow-up and data quality check, as well as the participants, will be
195 blinded regarding the shoe version distributed. The shoe code will be broken after completion of data
196 analysis.

197

198 **Intervention**

199 The study shoes are prototypes and will be anonymized for the purpose of this trial. The sole of the
200 shoes will be customized so that the two running shoe prototypes will be exactly the same (same
201 midsole, same outsole, same upper), except for their cushioning properties which will differ by about

35%, while remaining within the range of the models available on the market (stiffness: ~53-97 N/mm). The differences in cushioning properties between shoe versions will be created by modifying the midsole material, i.e. chemistry, density, and therefore the hardness of the Ethylene Vinyl Acetate (EVA) foam. In order to provide accurate data on the technical specifications (i.e. shoe stiffness) of each prototype, a set of 40 shoes (10 pairs per condition) will be tested for stiffness properties by the manufacturer according to a standardized protocol (Impact test: ASTM1614, Procedure A).[38]

Data collection

Baseline questionnaire

During the online registration process, the participants have to fill in a baseline questionnaire to report information regarding running experience, training habits, recent running competitions performed and injury history. A standardised questionnaire concerning the risk of sports participation must also be completed by the volunteers (Supplementary file 1). Every participant responding positively to any of the symptom-based questions or presenting more than one cardiovascular risk factor will be invited for a clearance check by a sports medical doctor prior to the test.

Biomechanical testing

The biomechanical running analysis will be performed on an instrumented treadmill (M-Gait, Motekforce Link Amsterdam, The Netherlands) in the randomly allocated study shoes. The test (10 minutes) consists of a 5-minute warm-up followed by a 5-minute run at the self-declared preferred (habitual) running speed. Two records of 45 seconds will be obtained at a sampling rate of 1 kHz over the last 2 minutes of the test. No data will be recorded during the first 8 minutes, which was shown to be enough time to provoke short-term adaptations of running style with respect to the shoe type.[25, 39] The main biomechanical variables of interest are presented in table 1.

Table 1: Biomechanical variables of interest.

Variable	Abbreviation	Unit	Normalization
----------	--------------	------	---------------

Step frequency	SF	[Steps.min ⁻¹]	/
Contact time	CT	[ms]	/
Flight time	FT	[ms]	/
Duty factor	DF	[%]	/
Step length	SL	[m]	[%LL]
Vertical Impact Peak Force	VIPF	[N]	[N.kg ⁻¹]
Peak Vertical Force	PVF	[N]	[N.kg ⁻¹]
Vertical Instantaneous Loading Rate	VILR	[N.s ⁻¹]	[N.kg ⁻¹ .s ⁻¹]
Vertical Average Loading Rate	VALR	[N.s ⁻¹]	[N.kg ⁻¹ .s ⁻¹]
Peak Power	PP	[W]	[W.kg ⁻¹]
Time to Peak Force	TPF	[ms]	/
Leg stiffness	Kleg	(kN/m)	/
Vertical stiffness	Kvert	(kN/m)	/

N: Newton, min: minute, ms: millisecond, m: meter, LL: leg length, kg: kilogram, W: Watt.

Anthropometric measures

Body mass and height of each participant will be measured barefoot and in running clothes before the treadmill running test. Also, the participants will have to report their body mass on a monthly basis onto their TIPPS account. Pop-up windows will inform the participants when an update is needed. In clinical settings, leg length is usually assessed as the measure between the anterior superior iliac spine and the medial malleolus (supine position), and is referred to as the “direct” clinical method.[40] The measurements will be performed on both legs and the average value will be used for the normalisation of step length. Additionally, the distance between the greater trochanter and the ground will be measured (standing position) to assess leg stiffness.[41] Body composition will be evaluated by bioelectrical impedance analysis (Tanita SC-240 MA). The proportion of fat mass will be included in the analyses as a potential confounder for the association between body mass and injury risk.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

242 *Data on exposure*

243 Data on running practice will be collected using the TIPPS system.[32, 42] Required information in
244 the sport activity report includes the type of activity, context, duration, subjectively perceived
245 intensity, distance, shoe pair used, running surface (hard or soft), and whether the participant had
246 experienced any pain during the session forcing him/her to reduce practice volume or intensity, or to
247 interrupt the practice. Session intensity is determined using the Borg’s rating of perceived exertion
248 scale, a subjective 10-point scale.[43]

250 *Data on outcome*

251 The primary outcome is the first RRI occurring during the follow-up. A consensus definition of RRI
252 in recreational runners has been recently published.[44] The definition of RRI is a “running-related
253 (training or competition) musculoskeletal pain in the lower limbs that causes a restriction on or
254 stoppage of running (distance, speed, duration, or training) for at least 7 days or 3 consecutive
255 scheduled training sessions, or that requires the runner to consult a physician or other health
256 professional.”

257 In previous studies, an RRI was defined as “any physical pain located at the lower limbs or lower
258 back region, sustained during or as a result of running practice and impeding planned running activity
259 for at least 1 day” (time-loss definition).[15, 32, 36, 37, 42] All painful episodes reported by the
260 participants during the follow-up will be assessed by a member of the research team according to each
261 of the two definitions presented above. The consensus definition will be considered as the reference,
262 while a sensitivity analysis will reveal if the results would be impacted when using the former
263 definition of RRI.

264 Similarly to uploading a training session or competition, the TIPPS provides a complete yet easy to
265 fill in questionnaire when reporting an injury. Information regarding the following is required: injury
266 date, context, sports discipline, injury mechanism (acute or progressive), anatomical location, type of
267 injury, description (free text field) and estimated return date. RRIs will be classified according to the

Orchard Sports Injury Classification System version 10 (OSICS-10).[45] Injury severity will be measured in days of modified or interrupted training.

Follow-up

Given that the participants are required to practice running at least once a week, individual e-mail reminders will be sent to the participants who do not provide the system with any data for the preceding week. Personal phone calls will be made if the participants do not react to the e-mail reminders and if the reported information in either the training log or on the injury form is found to be inconsistent.

Participants reporting any injury will be systematically contacted by one of the investigators to verify completeness and coherence of the reported data, and to check if the episode qualifies as an RRI (as defined above). Participants who do not complete their entire running calendar with weekly information will be contacted by one of the investigators to ensure that a RRI is not the reason for non-compliance or dropping out. The intervention period will last six months, allowing enough time for the participants to cover a large distance with the study shoes.

Sample size

A sample size calculation for Cox regression was used to determine the number of participants needed for the primary hypothesis of the study. With an alpha of 0.05 and a power of 80%, an average injury rate of 30%,[15, 36, 37] an expected hazard rate ratio (HR)=1.50 between groups, 50% of participants randomised to each shoe group and an expected drop-out rate of 20%, the total number of participants required is 802.

Statistical analysis

Descriptive data for the personal, anthropometric, biomechanical and training-related characteristics will be presented as count and percentage for dichotomous variables, and as mean and standard deviation, or as median and range, respectively, for normally and non-normally distributed continuous

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

variables. Average sport-related characteristics will be computed for each participant over their specific period of observation. Shock absorption properties (stiffness, N/mm) of the two types of shoes will be compared using a Student's t test.

Cox proportional hazards regressions will be used to compute the hazard rates in the exposure groups, using first-time injury as the primary outcome. Date of inclusion (baseline evaluation date) and date of injury or of censoring will be basic data used to calculate the time at risk, which is expressed in hours spent running and defined as the time-scale.[35] A participant will be right-censored if injury unrelated to running or severe disease caused a modification of the running plan, or at the end of follow-up. Reasons for right-censoring will be reported. The assumption of proportional hazards will be evaluated by log-minus-log plots.

Unadjusted Cox regressions will be performed to present the crude estimates of HRs for shoe model, body mass and other potential risk factors such as running biomechanics variables (see table 1) and training-related characteristics. Body mass is an exposure that can change over time (time-dependent covariate). This means that each participant could move between exposure states continuously (every month in our study). A delayed entry will be used in the unadjusted Cox regression model for body mass.[46]

Subsequently, the variables with a P value <0.200 will be included in the adjusted Cox regression analysis to determine whether shoe cushioning and/or body mass are associated with injury risk, controlling for potential confounders. The recommendation for using at least 10 injuries per predictor variable included in the Cox regression analysis will be strictly followed.[47]

Finally, to investigate if the effect of shoe cushioning on RRI risk is modified by body mass, a stratified analysis will be performed using the median value of body mass as cut-off. HRs and their 95% confidence intervals (CI) will be determined within each stratum.[48] All analyses will be performed using STATA/SE version 14.

DISCUSSION

It is common belief that shoe cushioning technology protects the runner against harmful consequences of repetitive high-load impacts. Therefore, heavier runners are generally advised to use footwear with adapted shock absorption properties. Surprisingly, few studies have investigated the impact of shoe cushioning on injury risk.[32, 33] These studies did not provide any evidence on the beneficial effect of increased shock absorption properties. However, none of them included anthropometric measures in their analyses. Also, one study compared different types of insoles added in the shoes,[33] while the other compared two versions of a standard running shoe with a limited difference in midsole hardness (~15%).[32] Other study limitations such as the sample size ($n < 250$)[32] or the study population (Royal Air Force recruits)[33] suggest that these results should be interpreted with caution. The evidence on the association between running shoe cushioning and RRI is still poor and inconclusive. One of the main reasons is the practical constraint of investigations trying to combine biomechanical analyses with a long-term prospective follow-up in a large number of runners.[11] This study is the first randomised controlled trial investigating the influence of shoe cushioning on RRI risk including an evaluation of running technique in all participants. The results will provide information on the real benefits provided by additional cushioning, as well as on the mechanisms that might explain any potential preventive effect.

337

338 ETHICS AND DISSEMINATION

This study will be conducted in accordance with the Declaration of Helsinki and the Medical Research Involving Human Subjects Act. Also, the study protocol (Ref: 201701/02 v1.1) was approved by the National Ethics Committee for Research (www.cner.lu). Written informed consent will be obtained from all participants (Supplementary file 2). All collected data will be stored electronically using a coding system. This will ensure that the data is used in the strictest confidence and will not reveal the identity of the participants. Collected raw data will not be passed on to unauthorised third parties. Results presented or published in articles and reports will be depicted in general terms, to maintain participant anonymity. Electronic data will be stored on a secure server in data files only accessible to the project leader and co-investigators of the project. A notification of this

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

study was sent to the National Data Protection Agency (CNPD). Study results will be submitted for publication in peer-reviewed journals and for presentation at international conferences. Furthermore, we aim to disseminate our results through popular specialised magazines and websites.

Contributors - LM, ND, AU and DT contributed to the study conception and study design. LM is the main investigator, wrote the article with input from other investigators, and will be responsible for the acquisition and analysis of the data. ND will be responsible for the shoe design, production and testing. LM and DT will be responsible for data interpretation and manuscript drafting. ND, AU and DT commented on the various versions of the study protocol. All authors approved the final manuscript.

Funding – This study is co-funded by Decathlon, Villeneuve d’Ascq, France and by the Ministry of Higher Education and Research of Luxembourg.

Competing interests – A research partnership agreement was signed between Decathlon and the Luxembourg Institute of Health (LIH). ND is employed at Decathlon SA. Decathlon will not be involved in the collection, management, analysis and interpretation of data. LM, DT and AU may not gain or lose financially from the results of the study in any way.

Ethics approval - All procedures were approved by the National Ethics Committee for Research (Ref: 201701/02 v1.1).

370 **REFERENCE LIST**

371 1 Kohl HW, 3rd, Craig CL, Lambert EV, et al. The pandemic of physical inactivity: global
372 action for public health. *Lancet*. 2012;**380**:294-305.

373 2 World Health Organization (WHO). Global status report on noncommunicable diseases
374 2014. . *Geneva: World Health Organization (WHO)*. 2014.

375 3 Hespanhol Junior LC, Pillay JD, van Mechelen W, et al. Meta-Analyses of the Effects of
376 Habitual Running on Indices of Health in Physically Inactive Adults. *Sports Med*.
377 2015;**45**:1455-68.

378 4 Lopes AD, Hespanhol Junior LC, Yeung SS, et al. What are the main running-related
379 musculoskeletal injuries? A Systematic Review. *Sports Med*. 2012;**42**:891-905.

380 5 Alentorn-Geli E, Samuelsson K, Musahl V, et al. The Association of Recreational and
381 Competitive Running With Hip and Knee Osteoarthritis: A Systematic Review and Meta-
382 Analysis. *J Orthop Sports Phys Ther*. 2017:1-36.

383 6 Koplan JP, Rothenberg RB, Jones EL. The natural history of exercise: a 10-yr follow-up of
384 a cohort of runners. *Med Sci Sports Exerc*. 1995;**27**:1180-4.

385 7 Hespanhol Junior LC, van Mechelen W, Postuma E, et al. Health and economic burden of
386 running-related injuries in runners training for an event: A prospective cohort study. *Scand J*
387 *Med Sci Sports*. 2016;**26**:1091-9.

388 8 Hespanhol Junior LC, van Mechelen W, Verhagen E. Health and Economic Burden of
389 Running-Related Injuries in Dutch Trailrunners: A Prospective Cohort Study. *Sports Med*.
390 2016.

391 9 van Gent RN, Siem D, van Middelkoop M, et al. Incidence and determinants of lower
392 extremity running injuries in long distance runners: a systematic review. *Br J Sports Med*.
393 2007;**41**:469-80; discussion 80.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

394 10 Knapik JJ, Jones BH, Steelman RA. Physical training in boots and running shoes: a
395 historical comparison of injury incidence in basic combat training. *Mil Med.* 2015;**180**:321-8.

396 11 Theisen D, Malisoux L, Gette P, et al. Footwear and running-related injuries –Running on
397 faith? *Sports Orthop Traumatol.* 2016;**32**:169-76.

398 12 Knapik JJ, Trone DW, Tchandja J, et al. Injury-reduction effectiveness of prescribing
399 running shoes on the basis of foot arch height: summary of military investigations. *J Orthop*
400 *Sports Phys Ther.* 2014;**44**:805-12.

401 13 Meeuwisse WH, Tyreman H, Hagel B, et al. A dynamic model of etiology in sport injury:
402 the recursive nature of risk and causation. *Clin J Sport Med.* 2007;**17**:215-9.

403 14 Hreljac A. Impact and overuse injuries in runners. *Med Sci Sports Exerc.* 2004;**36**:845-9.

404 15 Malisoux L, Nielsen RO, Urhausen A, et al. A step towards understanding the
405 mechanisms of running-related injuries. *J Sci Med Sport.* 2015;**18**:523-8.

406 16 van der Worp H, Vrielink JW, Bredeweg SW. Do runners who suffer injuries have higher
407 vertical ground reaction forces than those who remain injury-free? A systematic review and
408 meta-analysis. *Br J Sports Med.* 2016.

409 17 Davis IS, Bowser BJ, Mullineaux DR. Greater vertical impact loading in female runners
410 with medically diagnosed injuries: a prospective investigation. *Br J Sports Med.* 2015.

411 18 Barton CJ, Bonanno DR, Carr J, et al. Running retraining to treat lower limb injuries: a
412 mixed-methods study of current evidence synthesised with expert opinion. *Br J Sports Med.*
413 2016;**50**:513-26.

414 19 Diebal AR, Gregory R, Alitz C, et al. Forefoot running improves pain and disability
415 associated with chronic exertional compartment syndrome. *Am J Sports Med.* 2012;**40**:1060-
416 7.

- 20 Azevedo AP, Mezencio B, Amadio AC, et al. 16 Weeks of Progressive Barefoot Running Training Changes Impact Force and Muscle Activation in Habitual Shod Runners. *PLoS One*. 2016;**11**:e0167234.
- 21 Boyer ER, Derrick TR. Select injury-related variables are affected by stride length and foot strike style during running. *Am J Sports Med*. 2015;**43**:2310-7.
- 22 Luedke LE, Heiderscheit BC, Williams DS, et al. Influence of Step Rate on Shin Injury and Anterior Knee Pain in High School Runners. *Med Sci Sports Exerc*. 2016;**48**:1244-50.
- 23 Lorimer AV, Hume PA. Stiffness as a Risk Factor for Achilles Tendon Injury in Running Athletes. *Sports Med*. 2016;**46**:1921-38.
- 24 Baltich J, Maurer C, Nigg BM. Increased vertical impact forces and altered running mechanics with softer midsole shoes. *PLoS One*. 2015;**10**:e0125196.
- 25 Hardin EC, van den Bogert AJ, Hamill J. Kinematic adaptations during running: effects of footwear, surface, and duration. *Med Sci Sports Exerc*. 2004;**36**:838-44.
- 26 Nigg BM, Baltich J, Maurer C, et al. Shoe midsole hardness, sex and age effects on lower extremity kinematics during running. *J Biomech*. 2012;**45**:1692-7.
- 27 Shorten M, Mientjes M. The 'heel impact' force peak during running is neither 'heel' nor 'impact' and does not quantify shoe cushioning effects. *Footwear Science*. 2011;**3**:41-58.
- 28 Logan S, Hunter I, JT JTH, et al. Ground reaction force differences between running shoes, racing flats, and distance spikes in runners. *J Sports Sci Med*. 2010;**9**:147-53.
- 29 O'Leary K, Vorpahl KA, Heiderscheit B. Effect of cushioned insoles on impact forces during running. *J Am Podiatr Med Assoc*. 2008;**98**:36-41.
- 30 Kersting UG, Bruggemann GP. Midsole material-related force control during heel-toe running. *Res Sports Med*. 2006;**14**:1-17.
- 31 Nigg BM, Bahlsten HA, Luethi SM, et al. The influence of running velocity and midsole hardness on external impact forces in heel-toe running. *J Biomech*. 1987;**20**:951-9.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

442 32 Theisen D, Malisoux L, Genin J, et al. Influence of midsole hardness of standard
443 cushioned shoes on running-related injury risk. *Br J Sports Med.* 2014;**48**:371-6.

444 33 Withnall R, Eastaugh J, Freemantle N. Do shock absorbing insoles in recruits undertaking
445 high levels of physical activity reduce lower limb injury? A randomized controlled trial. *J R*
446 *Soc Med.* 2006;**99**:32-7.

447 34 Buist I, Bredeweg SW, Bessem B, et al. Incidence and risk factors of running-related
448 injuries during preparation for a 4-mile recreational running event. *Br J Sports Med.*
449 2010;**44**:598-604.

450 35 Nielsen RO, Malisoux L, Moller M, et al. Shedding Light on the Etiology of Sports
451 Injuries: A Look Behind the Scenes of Time-to-Event Analyses. *J Orthop Sports Phys Ther.*
452 2016:1-34.

453 36 Malisoux L, Chambon N, Delattre N, et al. Injury risk in runners using standard or motion
454 control shoes: a randomised controlled trial with participant and assessor blinding. *Br J*
455 *Sports Med.* 2016;**50**:481-7.

456 37 Malisoux L, Chambon N, Urhausen A, et al. Influence of the Heel-to-Toe Drop of
457 Standard Cushioned Running Shoes on Injury Risk in Leisure-Time Runners: A Randomized
458 Controlled Trial With 6-Month Follow-up. *Am J Sports Med.* 2016.

459 38 International A. ASTM F1614-99 Standard test method for shock attenuating properties of
460 materials systems for athletic footwear. *ASTM International.* 2006:West Conshohocken, PA,
461 USA.

462 39 Delattre N, Chambon N, Berton E, et al. Effect of time during a running session with
463 minimal footwear. *Comput Methods Biomech Biomed Engin.* 2013;**16 Suppl 1**:104-5.

464 40 Sabharwal S, Kumar A. Methods for assessing leg length discrepancy. *Clin Orthop Relat*
465 *Res.* 2008;**466**:2910-22.

- 466 41 Morin JB, Dalleau G, Kyrolainen H, et al. A simple method for measuring stiffness during
467 running. *J Appl Biomech*. 2005;**21**:167-80.
- 468 42 Malisoux L, Ramesh J, Mann R, et al. Can parallel use of different running shoes decrease
469 running-related injury risk? *Scand J Med Sci Sports*. 2015;**25**:110-5.
- 470 43 Foster C. Monitoring training in athletes with reference to overtraining syndrome. *Med Sci*
471 *Sports Exerc*. 1998;**30**:1164-8.
- 472 44 Yamato TP, Saragiotto BT, Lopes AD. A consensus definition of running-related injury in
473 recreational runners: a modified Delphi approach. *J Orthop Sports Phys Ther*. 2015;**45**:375-
474 80.
- 475 45 Rae K, Orchard J. The Orchard Sports Injury Classification System (OSICS) version 10.
476 *Clin J Sport Med*. 2007;**17**:201-4.
- 477 46 Nielsen RO, Parner ET, Nohr EA, et al. Excessive progression in weekly running distance
478 and risk of running-related injuries: an association which varies according to type of injury. *J*
479 *Orthop Sports Phys Ther*. 2014;**44**:739-47.
- 480 47 Peduzzi P, Concato J, Feinstein AR, et al. Importance of events per independent variable
481 in proportional hazards regression analysis. II. Accuracy and precision of regression
482 estimates. *J Clin Epidemiol*. 1995;**48**:1503-10.
- 483 48 Knol MJ, VanderWeele TJ. Recommendations for presenting analyses of effect
484 modification and interaction. *Int J Epidemiol*. 2012;**41**:514-20.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE LEGEND


Figure 1: Trial design.

SUPPLEMENTARY FILES

Supplementary file 1: Risk of sport participation form

Supplementary file 2: Informed consent

Figure 1: Trial Design



Trial design

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplementary file 4 : Questionnaire on risk of sport participation

- 1. Past medical history, have you had:
 - 1.1 Severe cardiac arrhythmia?
 - 1.2 Myocardial infarction?
 - 1.3 Heart surgery?
 - 1.4 Intracardiac catheter?
 - 1.5 Coronary angioplasty (dilatation by balloon, stenting)?
 - 1.6 Pacemaker or heart defibrillator?
 - 1.7 Cardiac insufficiency?
 - 1.8 Heart transplantation?
 - 1.9 Congenital heart defect?
- 2. Past and present complaints
 - 2.1 Chest pain / discomfort during physical exertion?
 - 2.2 Dyspnea (uncommon breathlessness)?
 - 2.3 Dizziness/unconsciousness?
 - 2.4 Palpitations, tachycardia, pulse irregularities?
 - 2.5 Intake of any cardiac drugs?
- 3. Other disorders
 - 3.1 Muscular or articular complaints?
 - 3.2 Other drugs?
 - 3.3 Insecurity during physical exertion?
 - 3.4 For females: Pregnancy?
- 4. Cardiovascular risk factors
 - 4.1 Are you male over 45 years?
 - 4.2 Are you female over 55 years or you have had a hysterectomy or you are menopausal?
 - 4.3 Smoker (active / in the past 10 years)?
 - 4.4 Your blood pressure is over 140/90 mmHg or you take antihypertensive drugs?
 - 4.5 Your cholesterol level is over 240 mg/dl?
 - 4.6 Myocardial infarction, stroke, marfan disease or sudden cardiac death in the family ? (Father resp. brother before age of 55 years/ Mother resp. sister before 65 years)?
 - 4.7 You are diabetic or you take antidiabetic drugs?
 - 4.8 Sports activity less than 90 min/week?
 - 4.9 You have a Body Mass Index (BMI) over 30 ?

Free Informed Consent

Title:

Institution:

Project manager:

Research assistant:

Head of unit:

1. I declare to have read the above-described information and accept to voluntarily participate in the study "Effects of bodyweight and shoe cushioning on injury risk and running biomechanics: A randomised control trial" conducted by the SMRL.
2. I accept that my data shall be used and communicated to the commercial partner for strictly scientific purposes once it has been pseudonymised (coded).
3. I received a copy of the present signed informed consent document, as well as the general information intended for athlete participants. I received a clear description of the purpose and the nature of the study and I am aware of what is expected of me as a participant in this study. I have had enough time and the opportunity to ask questions about the study; all my questions have been met with a satisfactory answer.
4. I am free to retire from the study at any time without justification. By doing so I will not suffer any material or moral damage.
5. I agree that the results of this study can be subject to public talks or scientific publication.
6. I voluntarily consent to participate in this study and I fully understand what kind of data will be gathered during the study.
7. I preserve/abide the rights of access, deletion or modification of my personal data. Any personal information will be kept confidential and protected in agreement with the modified personal data protection act of August 2nd 2002. I can exercise that right via the project manager.

The responding signatory freely consents to participate in the above mentioned study

Name and First Name of the respondent:

Signature of the respondent:

Name and signature of the project manager:

Place and date: